

THE MARYLAND ENTOMOLOGIST

Insect and related-arthropod studies in the Mid-Atlantic region



Volume 6, Number 4

September 2016

MARYLAND ENTOMOLOGICAL SOCIETY

www.mdentsoc.org

Executive Committee:	President	Frederick Paras
	Vice President	Philip J. Kean
	Secretary	Richard H. Smith, Jr.
	Treasurer	Edgar A. Cohen, Jr.
	Historian	(vacant)
	Journal Editor	Eugene J. Scarpulla
	E-newsletter Editor	Aditi Dubey & Hanna Kahl

The Maryland Entomological Society (MES) was founded in November 1971, to promote the science of entomology in all its sub-disciplines; to provide a common meeting venue for professional and amateur entomologists residing in Maryland, the District of Columbia, and nearby areas; to issue a periodical and other publications dealing with entomology; and to facilitate the exchange of ideas and information through its meetings and publications. The MES was incorporated in April 1982 and is a 501(c)(3) non-profit, scientific organization.

The MES logo features an illustration of *Euphydryas phaëton* (Drury) (Lepidoptera: Nymphalidae), the Baltimore Checkerspot, with its generic name above and its specific epithet below (both in capital letters), all on a pale green field; all these are within a yellow ring double-bordered by red, bearing the message “● Maryland Entomological Society ● 1971 ●”. All of this is positioned above the Shield of the State of Maryland. In 1973, the Baltimore Checkerspot was named the official insect of the State of Maryland through the efforts of many MES members.

Membership in the MES is open to all persons interested in the study of entomology. All members receive the annual journal, *The Maryland Entomologist*, and the monthly e-newsletter, *Phaëton*. Institutions may subscribe to *The Maryland Entomologist* but may not become members.

Annual Dues:	Individual Membership	\$10.00
	Household Membership	\$15.00
	Junior Membership (full-time student)	\$5.00
	Institutional Subscription	\$10.00

Prospective members should send to the Treasurer full dues for the current MES year (October – September), along with their full name, address, telephone number, e-mail address, and entomological interests. Applications can be downloaded from the MES website: www.mdentsoc.org. Send remittances, payable to the Maryland Entomological Society, and any address changes to the Treasurer: Edgar J. Cohen, Jr., 5454 Marsh Hawk Way, Columbia, MD 21045-2246, edcohenfam@yahoo.com.

Back issues of *The Maryland Entomologist* and recent issues of the *Phaëton* are available to members, via the Journal Editor, Eugene J. Scarpulla, ejscarp@comcast.net. Please contact the Journal Editor for availability and cost.

Meetings are held on the third Friday of October, November, February, March, April and May at 8:00 p.m. in Room 4 of the Biological Sciences Building, University of Maryland Baltimore County (UMBC), or occasionally at another announced site.

Past Presidents of the MES

1971-1973	Austin P. Platt	1984-1985	Charles L. Staines, Jr.
1973-1974	Ronald W. Hodges	1985-1986	Thomas E. Wallenmaier
1974-1975	Douglas C. Ferguson	1986-1987	Eugene J. Gerberg
1975	Raymond B. Nagle	1987-1988	Austin P. Platt
1975-1977	William A. Andersen	1988-1989	Philip J. Kean
1977-1978	Robert T. Mitchell	1989-1990	Nathan Erwin
1978-1979	Elaine R. Hodges	1990-1991	Stephen J. Harrison
1979-1980	Richard H. Smith, Jr.	1991-1992	Thomas E. Wallenmaier
1980-1981	Timothy P. Karpetsky	1992-1993	Mary C. Fenton
1981-1982	John F. Carroll	1993-2013	Frederick Paras
1982-1983	Theodore L. Bissell	2013-2015	Timothy Foard & Frederick Paras
1983-1984	Robin G. Todd	2015-present	Frederick Paras

Editor's Note

This year's issue of *The Maryland Entomologist* features an in-depth article titled "Specialist Bees of the Mid-Atlantic: Host Plants and Habitat Conservation" by Jarrod Fowler. Jarrod is the Pollinator Conservation and Conservation Biological Control Specialist for New England and Northeast Regions at The Xerces Society and a Technical Service Provider at the United States Department of Agriculture - Natural Resources Conservation Service. He has performed horticulture and entomology in New England for 20 years and is proficient in science-based and sustainable insect habitat assessment, design, installation, maintenance, and monitoring. Jarrod leads extensive insect habitat restoration efforts with farmers and landowners in New England.

Eugene J. Scarpulla
Editor

Specialist Bees of the Mid-Atlantic: Host Plants and Habitat Conservation

Jarrod Fowler

*The Xerces Society for Invertebrate Conservation, 628 NE Broadway, Suite 200,
Portland, Oregon 97232, USA
jarrod@xerces.org; (508) 274-1094*

Abstract: Habitat conservation is performed in North America to support populations of managed and wild pollinators. The current recommended plant selections for Mid-Atlantic pollinator habitats primarily provide resources for common or generalist pollinators. However, such plants may not foster uncommon or rare Mid-Atlantic specialist pollinators, whose populations are susceptible to harm from anthropogenic threats. This manuscript presents the first catalog of native specialist bees and associated host plants for the Mid-Atlantic. Approximately 29% of Mid-Atlantic native bee species are pollen specialists. I cataloged 6 families, 25 genera, and 131 species of Mid-Atlantic native bees that restrict their pollen use to 28 families, 54 genera, and 396 possible species of host plants. Specialist bees are associated with non-graminoid forbs and non-coniferous woody plants in nearly all Mid-Atlantic terrestrial and wetland habitats. Vulnerable bee-plant associations requiring greater emphasis in research and restoration are identified and discussed.

Keywords: Apoidea, Anthophila, habitat restoration, Mid-Atlantic, oligolecty, pollen specialization

INTRODUCTION

Mid-Atlantic native-pollen specialist bees (oligoleges) have evolved a continuum of facultative or obligate associations with flowering host plants (Robertson 1925, Linsley and MacSwain 1958, Hurd et al. 1980, Cane and Sipes 2006). Oligoleges associate with one host plant family or a few related genera or species; monoleges specifically associate with a single host plant genus or species (Robertson 1925, Cane and Sipes 2006). Such associations can benefit both bee and flower species by improving foraging effectiveness and efficiency, pollen digestibility, and pollination rates. However, restricted associations can be susceptible to harm due to habitat degradation, fragmentation, and loss, or to phenological mismatch (Minckley et al. 1994, Bartomeus et al. 2013, Rafferty et al. 2015). Consequently, anthropogenic threats in the Mid-Atlantic potentially imperil native specialist bee species and their endemic, indigenous host plant species with population declines and extinctions through loss of plant diversity (Radeloff et al. 2005).

Pollen associations and preferences for many Mid-Atlantic native specialist bees can be derived from many published and unpublished field observations and specimen records (Lovell 1918; Robertson 1926, 1929; Mitchell 1960, 1962). Here, I organize bee-plant observations and records to catalog Mid-Atlantic native host plants for native specialist

bees. This catalog provides opportunities to augment current lists of plant selections for pollinator habitat conservation and enhancement with recommended host plants for specialist bees. Such recommendations can be integrated into pollinator habitat practices, and are broadly relevant to the Mid-Atlantic and the adjacent regions in Canada and the Northeastern United States.

Field-site Description

This research includes pollination records from the Mid-Atlantic United States for six states: Delaware, Maryland, New Jersey, Pennsylvania, Virginia, and West Virginia. New York is often included in definitions of what comprises the Mid-Atlantic, but has been excluded from consideration. The combination of “Specialist Bees of the Northeast: Host Plants and Habitat Conservation” (Fowler 2016) and the present manuscript will encompass most bee-plant associations found in New York. The District of Columbia (DC) has been similarly excluded from consideration. Urbanized DC is geographically minute compared to surrounding Maryland and Virginia.

The Mid-Atlantic is an ecologically and topographically diverse region within the temperate broadleaf and mixed forest biome. Topography ranges from western low mountains and plateaus, with elevations up to 1,750 m (5741 ft), to outwash plains of unconsolidated sediments that create a large coastal plain which contains extensive marshes, barrier islands, and dune systems (Omernik 1987; Jones et al. 1997). Overall, humid cold temperate and warm continental climates with moderate precipitation and moist yet well-drained soils foster species-dense and species-rich communities of native, endemic, exotic, and hybrid terrestrial, wetland, and aquatic plant species (Adamson et al. 2015).

METHODS

Here specialist bees are defined as those that restrict pollen foraging from only one host plant family (Robertson 1925, Cane and Sipes 2006). In reality, most specialist bee records are associated with one host plant genus or species. Specialist bees that are indigenous to Delaware, Maryland, New Jersey, Pennsylvania, Virginia, and West Virginia without human intervention were defined as Mid-Atlantic natives. Specimen records of approximately 450 species of native bees captured or observed on flowers of host plants were compiled from online sources (Hilty 2012, Ascher and Pickering 2016), peer reviewed articles (Robertson 1926, 1929; LaBerge 1969, 1971, 1973, 1977, 1980, 1985, 1986, 1989; LaBerge and Bouseman 1970; LaBerge and Ribble 1972, 1975; Bouseman and LaBerge 1978), technical bulletins (Mitchell 1960, 1962; LaBerge 1967; Krombein et al. 1979), and personal communications with contemporary bee researchers. Information concerning conservation statuses, habitats, native ranges, nesting organizations, monthly phenologies, numbers of records per state, and taxonomies was collected in addition to host plant associations for each bee species. Cleptoparasitic bees were not included in our study.

Bee-plant records were compared with state-level Mid-Atlantic plant distributions from the United States Department of Agriculture - Natural Resources Conservation Service PLANTS Database (USDA-NRCS 2016). According to plant distribution source

definitions, native species are historically indigenous prior to European colonization and present species have been documented as uncultivated native or naturalized populations (USDA-NRCS 2016). This study includes only host plant genera and species that are native and documented as present in at least one of the six Mid-Atlantic states. If there are records of specialist bee species collected from Mid-Atlantic native plant genera or species, even if those collections came from outside the Mid-Atlantic, then such plant taxa are listed. Because plant taxonomy has recently undergone major changes, many names have been updated compared to those published in the original literature of bee-plant associations. Information concerning plant taxonomy, morphological groupings, native ranges, and habitat types (obligate–facultative–upland) was compiled for each host plant species. As above, such data were analyzed and summarized to provide recommendations for targeted conservation and management in the Mid-Atlantic and nearby regions.

RESULTS

One hundred thirty-one species, or 29% of approximately 450 Mid-Atlantic native bee species, were pollen specialists. Six families, 25 genera, and 131 species of Mid-Atlantic native bees and 28 families, 54 genera, and 396 possible species of host plants were cataloged. Specialist bees are alphabetically tabulated with associated host plant families and genera in Table 1. Conversely, host plant families and genera with hosted bee families and species are summarized in Table 2. The most recurrent host plant family association among specialist bee species was Asteraceae, 58 bee spp. The most recurrent host plant genera associations among specialist bee species were *Salix* L. (willows), 11 bee spp.; *Solidago* L. (goldenrods), 11 bee spp.; *Vaccinium* L. (blueberries), 8 bee spp.; *Helianthus* L. (sunflowers), 7 bee spp.; and *Symphyotrichum* Nees (asters), 7 bee spp.

Six families, 25 genera, and 131 species of Mid-Atlantic native specialist bees were cataloged. Life history and taxonomic information about Mid-Atlantic native specialist bees are presented in Supplemental Table 3. The bee family with the most specialists in the Mid-Atlantic was Andrenidae (mining bees), 70 spp., and the bee genus with most species was *Andrena* Fabricius, 47 spp. Of the 131 bee species, 66 were considered rare, while 49 were uncommon, and 15 were common. All bee species presented are solitary nesters and none are known to be eusocial, although some might be communal (e.g., *Perdita* Smith [Andrenidae]) or aggregate nesters (e.g., *Colletes* Latreille [Colletidae]). Additionally, most cataloged bees are soil nesters with the exception of 13 Megachilidae (leafcutter, mason, and resin bees), which likely nest in hollow/pithy stems or holes in wood. Overall numbers of specialist bee species active per family per month are presented in Figure 1. Specialist bee adults were effectively inactive during December, January, and February in the Mid-Atlantic, remaining in their natal nests as pre-pupae or as adults in some early-spring emerging species.

Associations with non-graminoid (non-grass or non-grass-like) forbs and non-coniferous woody plants were clear for all Mid-Atlantic specialist bee species. Twenty-eight families, 54 genera, and 396 possible species of native host plants were cataloged. Life history and taxonomic information about Mid-Atlantic native host plants are presented in Supplemental Table 4. Overall, the host plant families with the greatest number of species were Asteraceae (140 spp.), Ericaceae (33 spp.), Brassicaceae (26 spp.), and Violaceae (26 spp.). The host plant genera with the greatest number of species were

Table 1. Mid-Atlantic native specialist bee families and species with their associated host plant families and genera. Bee families and member genera and species with associated host plants are presented in ascending alphabetical order. Consult Supplementary Tables 3 and 4 for additional information.

Bee Family	Bee Species	Host Plant Family	Host Plant Genus
Andrenidae	<i>Andrena accepta</i>	Asteraceae	<i>Helianthus</i>
	<i>Andrena aliciae</i>	Asteraceae	<i>Helianthus</i>
	<i>Andrena andrenoides</i>	Salicaceae	<i>Salix</i>
	<i>Andrena arabis</i>	Brassicaceae	<i>Arabis</i>
			<i>Cardamine</i>
	<i>Andrena asteris</i>	Asteraceae	<i>Solidago</i>
			<i>Symphyotrichum</i>
	<i>Andrena asteroides</i>	Asteraceae	<i>Symphyotrichum</i>
	<i>Andrena bisalicens</i>	Salicaceae	<i>Salix</i>
	<i>Andrena braccata</i>	Asteraceae	<i>Euthamia</i>
			<i>Solidago</i>
	<i>Andrena bradleyi</i>	Ericaceae	<i>Gaylussacia</i>
			<i>Vaccinium</i>
	<i>Andrena canadensis</i>	Asteraceae	<i>Solidago</i>
	<i>Andrena carolina</i>	Ericaceae	<i>Vaccinium</i>
	<i>Andrena chromotricha</i>	Asteraceae	.
	<i>Andrena clarkella</i>	Salicaceae	<i>Salix</i>
	<i>Andrena cornelli</i>	Ericaceae	<i>Rhododendron</i>
	<i>Andrena distans</i>	Geraniaceae	<i>Geranium</i>
	<i>Andrena duplicata</i>	Asteraceae	.
	<i>Andrena erigeniae</i>	Portulacaceae	<i>Claytonia</i>
	<i>Andrena erythrogaster</i>	Salicaceae	<i>Salix</i>
	<i>Andrena erythronii</i>	Liliaceae	<i>Erythronium</i>
	<i>Andrena fragilis</i>	Cornaceae	<i>Cornus</i>
	<i>Andrena frigida</i>	Salicaceae	<i>Salix</i>
	<i>Andrena fulvipennis</i>	Asteraceae	<i>Chrysopsis</i>
			<i>Heterotheca</i>
			<i>Pityopsis</i>
	<i>Andrena gardineri</i>	Asteraceae	<i>Packera</i>
	<i>Andrena geranii</i>	Hydrophyllaceae	<i>Hydrophyllum</i>
	<i>Andrena helianthi</i>	Asteraceae	<i>Helianthus</i>
	<i>Andrena hirticincta</i>	Asteraceae	<i>Euthamia</i>
			<i>Solidago</i>
	<i>Andrena illinoensis</i>	Salicaceae	<i>Salix</i>
	<i>Andrena integra</i>	Cornaceae	<i>Cornus</i>
	<i>Andrena krigiana</i>	Asteraceae	<i>Krigia</i>
	<i>Andrena lamelliterga</i>	Hydrophyllaceae	<i>Phacelia</i>
	<i>Andrena macoupinensis</i>	Salicaceae	<i>Salix</i>
	<i>Andrena mariae</i>	Salicaceae	<i>Salix</i>
	<i>Andrena melanochroa</i>	Rosaceae	<i>Fragaria</i>
	<i>Andrena nida</i>	Salicaceae	<i>Salix</i>
	<i>Andrena nigrae</i>	Salicaceae	<i>Salix</i>
	<i>Andrena nubecula</i>	Asteraceae	<i>Solidago</i>
			<i>Symphyotrichum</i>
	<i>Andrena phaceliae</i>	Hydrophyllaceae	<i>Phacelia</i>
	<i>Andrena placata</i>	Asteraceae	<i>Solidago</i>
			<i>Symphyotrichum</i>

Bee Family	Bee Species	Host Plant Family	Host Plant Genus
	<i>Andrena platyparia</i>	Cornaceae	<i>Cornus</i>
	<i>Andrena polemonii</i>	Polemoniaceae	<i>Polemonium</i>
	<i>Andrena rudbeckiae</i>	Asteraceae	<i>Rudbeckia</i>
	<i>Andrena salictaria</i>	Salicaceae	<i>Salix</i>
	<i>Andrena simplex</i>	Asteraceae	<i>Solidago</i> <i>Symphyotrichum</i>
	<i>Andrena uvulariae</i>	Liliaceae	<i>Uvularia</i>
	<i>Andrena violae</i>	Violaceae	<i>Viola</i>
	<i>Andrena ziziae</i>	Apiaceae	<i>Zizia</i>
	<i>Andrena ziziaeformis</i>	Rosaceae	<i>Potentilla, Waldsteinia</i>
	<i>Calliopsis nebraskensis</i>	Verbenaceae	<i>Verbena</i>
	<i>Panurginus atramontensis</i>	Ericaceae	<i>Vaccinium</i>
	<i>Panurginus potentillae</i>	Rosaceae	<i>Potentilla</i>
	<i>Perdita bequaerti</i>	Asteraceae	<i>Helianthus</i>
	<i>Perdita bishoppi</i>	Asteraceae	<i>Chrysopsis</i> <i>Heterotheca</i> <i>Pityopsis</i>
	<i>Perdita boltoniae</i>	Asteraceae	<i>Chrysopsis</i> <i>Heterotheca</i> <i>Pityopsis</i>
	<i>Perdita consobrina</i>	Asteraceae	.
	<i>Perdita gerardiae</i>	Scrophulariaceae	<i>Agalinis</i>
	<i>Perdita gerhardi</i>	Lamiaceae	<i>Monarda</i>
	<i>Perdita halictoides</i>	Solanaceae	<i>Physalis</i>
	<i>Perdita octomaculata</i>	Asteraceae	<i>Solidago</i>
	<i>Perdita swenki</i>	Asteraceae	.
	<i>Protandrena abdominalis</i>	Lamiaceae	<i>Monarda</i>
	<i>Pseudopanurgus aestivalis</i>	Asteraceae	<i>Solidago</i> <i>Symphyotrichum</i>
	<i>Pseudopanurgus albitarsis</i>	Asteraceae	.
	<i>Pseudopanurgus andrenoides</i>	Asteraceae	.
	<i>Pseudopanurgus compositarum</i>	Asteraceae	.
	<i>Pseudopanurgus illinoiensis</i>	Asteraceae	.
	<i>Pseudopanurgus labrosiformis</i>	Asteraceae	.
	<i>Pseudopanurgus labrosus</i>	Asteraceae	.
	<i>Pseudopanurgus pauper</i>	Rhamnaceae	<i>Ceanothus</i>
	<i>Pseudopanurgus rugosus</i>	Asteraceae	.
	<i>Pseudopanurgus virginicus</i>	Rubiaceae	<i>Houstonia</i>
Apidae	<i>Anthophorula micheneri</i>	Scrophulariaceae	<i>Agalinis</i>
	<i>Cemolobus ipomoeae</i>	Convolvulaceae	<i>Ipomoea</i>
	<i>Florilegus condignus</i>	Pontederiaceae	<i>Pontederia</i>
	<i>Habropoda laboriosa</i>	Ericaceae	<i>Cercis</i> <i>Vaccinium</i>
	<i>Melissodes agilis</i>	Asteraceae	<i>Helianthus</i>
	<i>Melissodes apicatus</i>	Pontederiaceae	<i>Pontederia</i>
	<i>Melissodes boltoniae</i>	Asteraceae	.
	<i>Melissodes denticulatus</i>	Asteraceae	<i>Vernonia</i>
	<i>Melissodes dentiventris</i>	Asteraceae	.
	<i>Melissodes desponsus</i>	Asteraceae	<i>Cirsium</i>
	<i>Melissodes druriellus</i>	Asteraceae	.
	<i>Melissodes fimbriatus</i>	Onagraceae	<i>Oenothera</i>
	<i>Melissodes fumosus</i>	Asteraceae	.
	<i>Melissodes illatus</i>	Asteraceae	.
	<i>Melissodes niveus</i>	Asteraceae	.

Bee Family	Bee Species	Host Plant Family	Host Plant Genus
	<i>Melissodes subillatus</i>	Asteraceae	.
	<i>Melissodes trinodis</i>	Asteraceae	.
	<i>Melitoma taurea</i>	Convolvulaceae	<i>Ipomoea</i>
	<i>Ptilothrix bombiformis</i>	Malvaceae	<i>Hibiscus</i>
	<i>Svastra compta</i>	Onagraceae	<i>Oenothera</i>
	<i>Svastra obliqua</i>	Asteraceae	.
	<i>Svastra petulca</i>	Asteraceae	.
Colletidae	<i>Colletes aestivalis</i>	Saxifragaceae	<i>Heuchera</i>
	<i>Colletes banksi</i>	Aquifoliaceae	<i>Ilex</i>
	<i>Colletes brevicornis</i>	Campanulaceae	<i>Campanula</i> <i>Triodanis</i>
	<i>Colletes compactus</i>	Asteraceae	.
	<i>Colletes latitarsis</i>	Solanaceae	<i>Physalis</i>
	<i>Colletes productus</i>	Ericaceae	<i>Lyonia</i> <i>Vaccinium</i>
	<i>Colletes simulans</i>	Asteraceae	<i>Euthamia</i> <i>Solidago</i> <i>Symphytotrichum</i>
	<i>Colletes solidaginis</i>	Asteraceae	<i>Solidago</i>
	<i>Colletes speculiferus</i>	Asteraceae	.
	<i>Colletes thysanellae</i>	Asteraceae	.
	<i>Colletes validus</i>	Ericaceae	.
	<i>Colletes willistoni</i>	Solanaceae	<i>Physalis</i>
Halictidae	<i>Dieunomia heteropoda</i>	Asteraceae	<i>Bidens</i> <i>Helianthus</i>
	<i>Dieunomia nevadensis</i>	Asteraceae	.
	<i>Dufourea monardae</i>	Lamiaceae	<i>Monarda</i>
	<i>Dufourea novaeangliae</i>	Pontederiaceae	<i>Pontederia</i>
	<i>Lasioglossum lustrans</i>	Asteraceae	<i>Pyrrhopappus</i>
	<i>Lasioglossum oenotherae</i>	Onagraceae	<i>Oenothera</i>
	<i>Lasioglossum pectinatum</i>	Solanaceae	<i>Physalis</i>
Megachilidae	<i>Hoplitis simplex</i>	Hydrophyllaceae	<i>Nemophila</i> <i>Phacelia</i>
	<i>Megachile inimica</i>	Asteraceae	.
	<i>Megachile integra</i>	Fabaceae	<i>Galactia</i> <i>Strophostyles</i>
	<i>Megachile oenotherae</i>	Onagraceae	<i>Oenothera</i>
	<i>Megachile parallela</i>	Asteraceae	.
	<i>Megachile pugnata</i>	Asteraceae	<i>Helianthus</i>
	<i>Osmia chalybea</i>	Asteraceae	<i>Cirsium</i>
	<i>Osmia distincta</i>	Scrophulariaceae	<i>Penstemon</i>
	<i>Osmia texana</i>	Asteraceae	<i>Cirsium</i>
	<i>Osmia virga</i>	Ericaceae	<i>Vaccinium</i>
	<i>Paranthidium jugatorium</i>	Asteraceae	<i>Helianthus</i>
	<i>Trachusa dorsalis</i>	Fabaceae	<i>Strophostyles</i>
Melittidae	<i>Macropis ciliata</i>	Primulaceae	<i>Lysimachia</i>
	<i>Macropis nuda</i>	Primulaceae	<i>Lysimachia</i>
	<i>Macropis patellata</i>	Primulaceae	<i>Lysimachia</i>
	<i>Macropis steironematis</i>	Primulaceae	<i>Lysimachia</i>
	<i>Melitta americana</i>	Ericaceae	<i>Vaccinium</i>
	<i>Melitta eickworti</i>	Ericaceae	<i>Vaccinium</i>
	<i>Melitta melittoides</i>	Ericaceae	<i>Lyonia</i>

Table 2. Mid-Atlantic native plant families and genera with their hosted bee families and species. Plant families and member genera with hosted bees are presented in ascending alphabetical order. Consult Supplementary Tables 3 and 4 for additional information.

Host Plant Family	Host Plant Genus	Bee Family	Bee Species
Apiaceae	<i>Zizia</i>	Andrenidae	<i>Andrena ziziae</i>
Aquifoliaceae	<i>Ilex</i>	Colletidae	<i>Colletes banksi</i>
Asteraceae	.	Andrenidae	<i>Andrena chromotricha</i>
			<i>Andrena duplicata</i>
			<i>Perdita consobrina</i>
			<i>Perdita swenki</i>
			<i>Pseudopanurgus albitarsis</i>
			<i>Pseudopanurgus andrenoides</i>
			<i>Pseudopanurgus compositarum</i>
			<i>Pseudopanurgus illinoiensis</i>
			<i>Pseudopanurgus labrosiformis</i>
			<i>Pseudopanurgus labrosus</i>
			<i>Pseudopanurgus rugosus</i>
		Apidae	<i>Melissodes boltoniae</i>
			<i>Melissodes dentiventris</i>
			<i>Melissodes druriellus</i>
			<i>Melissodes fumosus</i>
			<i>Melissodes illatus</i>
			<i>Melissodes niveus</i>
			<i>Melissodes subillatus</i>
			<i>Melissodes trinodis</i>
			<i>Svastra obliqua</i>
			<i>Svastra petulca</i>
		Colletidae	<i>Colletes compactus</i>
			<i>Colletes speculiferus</i>
			<i>Colletes thysanellae</i>
		Halictidae	<i>Dieunomia nevadensis</i>
		Megachilidae	<i>Megachile inimica</i>
			<i>Megachile parallela</i>
			<i>Megachile xylocopoides</i>
	<i>Bidens</i>	Halictidae	<i>Dieunomia heteropoda</i>
	<i>Chrysopsis</i>	Andrenidae	<i>Andrena fulvipennis</i>
			<i>Perdita bishoppi</i>
			<i>Perdita boltoniae</i>
	<i>Cirsium</i>	Apidae	<i>Melissodes desponsus</i>
		Megachilidae	<i>Osmia chalybea</i>
			<i>Osmia texana</i>
	<i>Euthamia</i>	Andrenidae	<i>Andrena braccata</i>
			<i>Andrena hirticincta</i>
	<i>Helianthus</i>	Colletidae	<i>Colletes simulans</i>
		Andrenidae	<i>Andrena accepta</i>
			<i>Andrena aliciae</i>
			<i>Andrena helianthi</i>
			<i>Perdita bequaerti</i>
		Apidae	<i>Melissodes agilis</i>
		Halictidae	<i>Dieunomia heteropoda</i>
		Megachilidae	<i>Megachile pugnata</i>

Host Plant Family	Host Plant Genus	Bee Family	Bee Species
			<i>Paranthidium jugatorium</i>
	<i>Heterotheca</i>	Andrenidae	<i>Andrena fulvipennis</i>
			<i>Perdita bishoppi</i>
			<i>Perdita boltoniae</i>
	<i>Krigia</i>	Andrenidae	<i>Andrena krigiana</i>
	<i>Packera</i>	Andrenidae	<i>Andrena gardineri</i>
	<i>Pityopsis</i>	Andrenidae	<i>Andrena fulvipennis</i>
			<i>Perdita bishoppi</i>
			<i>Perdita boltoniae</i>
	<i>Pyrrhopappus</i>	Halictidae	<i>Lasioglossum lustrans</i>
	<i>Rudbeckia</i>	Andrenidae	<i>Andrena rudbeckiae</i>
	<i>Solidago</i>	Andrenidae	<i>Andrena asteris</i>
			<i>Andrena braccata</i>
			<i>Andrena canadensis</i>
			<i>Andrena hirticincta</i>
			<i>Andrena nubecula</i>
			<i>Andrena placata</i>
			<i>Andrena simplex</i>
			<i>Perdita octomaculata</i>
			<i>Pseudopanurgus nebrascensis</i>
		Colletidae	<i>Colletes simulans</i>
			<i>Colletes solidaginis</i>
	<i>Symphyotrichum</i>	Andrenidae	<i>Andrena asteris</i>
			<i>Andrena asteroides</i>
			<i>Andrena nubecula</i>
			<i>Andrena placata</i>
			<i>Andrena simplex</i>
			<i>Pseudopanurgus nebrascensis</i>
		Colletidae	<i>Colletes simulans</i>
	<i>Vernonia</i>	Apidae	<i>Melissodes denticulatus</i>
Brassicaceae	<i>Arabis</i>	Andrenidae	<i>Andrena arabis</i>
	<i>Cardamine</i>	Andrenidae	<i>Andrena arabis</i>
Campanulaceae	<i>Campanula</i>	Colletidae	<i>Colletes brevicornis</i>
	<i>Triodanis</i>	Colletidae	<i>Colletes brevicornis</i>
Convolvulaceae	<i>Ipomoea</i>	Apidae	<i>Cemolobus ipomoeae</i>
			<i>Melitoma taurea</i>
Cornaceae	<i>Cornus</i>	Andrenidae	<i>Andrena fragilis</i>
			<i>Andrena integra</i>
			<i>Andrena platyparia</i>
Ericaceae	.	Colletidae	<i>Colletes validus</i>
	<i>Gaylussacia</i>	Andrenidae	<i>Andrena bradleyi</i>
	<i>Lyonia</i>	Colletidae	<i>Colletes productus</i>
		Melittidae	<i>Melitta melittoides</i>
	<i>Rhododendron</i>	Andrenidae	<i>Andrena cornelli</i>
	<i>Vaccinium</i>	Andrenidae	<i>Andrena bradleyi</i>
			<i>Andrena carolina</i>
			<i>Panurginus atramontensis</i>
		Apidae	<i>Habropoda laboriosa</i>
		Colletidae	<i>Colletes productus</i>
		Megachilidae	<i>Osmia virga</i>
		Melittidae	<i>Melitta americana</i>
			<i>Melitta eickworti</i>
Fabaceae	<i>Cercis</i>	Apidae	<i>Habropoda laboriosa</i>
	<i>Galactia</i>	Megachilidae	<i>Megachile integra</i>

Host Plant Family	Host Plant Genus	Bee Family	Bee Species
	<i>Strophostyles</i>	Megachilidae	<i>Megachile integra</i> <i>Trachusa dorsalis</i>
Geraniaceae	<i>Geranium</i>	Andrenidae	<i>Andrena distans</i>
Hydrophyllaceae	<i>Hydrophyllum</i>	Andrenidae	<i>Andrena geranii</i>
	<i>Nemophila</i>	Megachilidae	<i>Hoplitis simplex</i>
	<i>Phacelia</i>	Andrenidae	<i>Andrena lamelliterga</i> <i>Andrena phaceliae</i>
		Megachilidae	<i>Hoplitis simplex</i>
Lamiaceae	<i>Monarda</i>	Andrenidae	<i>Perdita gerhardi</i> <i>Protandrena abdominalis</i>
		Halictidae	<i>Dufourea monardae</i>
Liliaceae	<i>Erythronium</i>	Andrenidae	<i>Andrena erythronii</i>
	<i>Uvularia</i>	Andrenidae	<i>Andrena uvulariae</i>
Malvaceae	<i>Hibiscus</i>	Apidae	<i>Ptilothrix bombiformis</i>
Onagraceae	<i>Oenothera</i>	Apidae	<i>Melissodes fimbriatus</i> <i>Svastra compta</i>
		Halictidae	<i>Lasioglossum oenotherae</i>
		Megachilidae	<i>Megachile oenotherae</i>
Polemoniaceae	<i>Polemonium</i>	Andrenidae	<i>Andrena polemonii</i>
Pontederiaceae	<i>Pontederia</i>	Apidae	<i>Florilegus condignus</i> <i>Melissodes apicata</i>
		Halictidae	<i>Dufourea novaeangliae</i>
Portulacaceae	<i>Claytonia</i>	Andrenidae	<i>Andrena erigeniae</i>
Primulaceae	<i>Lysimachia</i>	Melittidae	<i>Macropis ciliata</i> <i>Macropis nuda</i> <i>Macropis patellata</i> <i>Macropis steironematis</i>
Rhamnaceae	<i>Ceanothus</i>	Andrenidae	<i>Pseudopanurgus pauper</i>
Rosaceae	<i>Fragaria</i>	Andrenidae	<i>Andrena melanochroa</i>
	<i>Potentilla</i>	Andrenidae	<i>Andrena ziziaeformis</i> <i>Panurginus potentillae</i>
	<i>Waldsteinia</i>	Andrenidae	<i>Andrena ziziaeformis</i>
Rubiaceae	<i>Houstonia</i>	Andrenidae	<i>Pseudopanurgus virginicus</i>
Salicaceae	<i>Salix</i>	Andrenidae	<i>Andrena andrenoides</i> <i>Andrena bisalicens</i> <i>Andrena clarkella</i> <i>Andrena erythrogaster</i> <i>Andrena frigida</i> <i>Andrena illinoensis</i> <i>Andrena macoupinensis</i> <i>Andrena mariae</i> <i>Andrena nida</i> <i>Andrena nigrae</i> <i>Andrena salictaria</i>
Saxifragaceae	<i>Heuchera</i>	Colletidae	<i>Colletes aestivalis</i>
Scrophulariaceae	<i>Agalinis</i>	Andrenidae	<i>Perdita gerardiae</i>
		Apidae	<i>Anthophorula micheneri</i>
	<i>Penstemon</i>	Megachilidae	<i>Osmia distincta</i>
Solanaceae	<i>Physalis</i>	Andrenidae	<i>Perdita halictoides</i>
		Colletidae	<i>Colletes latitarsis</i> <i>Colletes willistoni</i>
		Halictidae	<i>Lasioglossum pectinatum</i>
Verbenaceae	<i>Verbena</i>	Andrenidae	<i>Calliopsis nebraskensis</i>
Violaceae	<i>Viola</i>	Andrenidae	<i>Andrena violae</i>




Figure 1. Numbers of Mid-Atlantic specialist bee species active per family per month.

Specialist bee adults were inactive during December, January, and February in the Mid-Atlantic. Overall numbers of species per month were: July (75 spp.), June (73 spp.), August (69 spp.), May (64 spp.), September (58 spp.), April (46 spp.), October (30 spp.), March (22 spp.), and November (8 spp.).

Solidago (35 spp.); *Symphyotrichum* (32 spp.); *Viola* L., violets, (26 spp.); *Helianthus* (18 spp.), *Oenothera* L., evening primroses, (17 spp.), and *Salix* (16 spp.). Host plants were primarily “herbaceous, flowering dicots” (317 spp.) and secondarily “broad-leaf woody shrubs” (40 spp.), “broad-leaf woody shrub-trees” (29 spp.), “herbaceous, flowering monocots” (8 spp.), and “broad-leaf woody trees” (2 spp.).

Associations with terrestrial and wetland habitats were clear for all Mid-Atlantic specialist bee species. The host plant habitat types (266 spp.) were primarily “facultative-wetland” (51 spp.) and secondarily “facultative-upland” (49 spp.), “facultative” (45 spp.), “obligate” (40 spp.), and “facultative-facultative-upland” (25 spp.). Of the 396 possible species of host plants, 171 species were native in all six Mid-Atlantic states.

DISCUSSION

This manuscript presents the first catalog of specialist bees and associated host plants for the Mid-Atlantic United States. Specialist bees, associated host plants, and future work will be discussed in the following paragraphs. In the process, recommendations will be provided according to the results.

Bees

The distribution of pollen specialists among the bee families, genera, and subgenera of the Mid-Atlantic is not random. Family Melittidae (oil bees) showed clear preferences: all three listed species of *Melitta* Kirby were associated with Ericaceae, while all four catalogued species of *Macropis* Panzer were associated with *Lysimachia* L. (yellow loosestrifes). All 11 cataloged species of *Andrena* subgenus *Callandrena* sensu lato and four cataloged species of *Andrena* subgenus *Cnemidandrena* were associated with Asteraceae. Similarly, all 10 cataloged species of *Melissodes* Latreille (Apidae; long-horned bees) subgenus *Eumelissodes* were associated with Asteraceae. Moreover, all three catalogued species of *Andrena* subgenus *Gonandrena* associated with *Cornus* L. (dogwoods).

Notably, the dominant and speciose bees in the families Halictidae (sweat bees) and Megachilidae are nearly devoid of oligoleges. Social bumble bees, *Bombus* Latreille (Apidae), have no pollen specialists, and oligoleges are similarly absent in the common masked bee genus *Hylaeus* Fabricius (Colletidae). Generalist bees of polyphagous and social groups do not restrict pollen, but do have broad and clear preferences for flowers based on body size, shape, and tongue morphology.

Several bee species deserve explanation for their exclusion. The Squash Bee, *Peponapis pruinosa* (Say), and its less common close relative *Xenoglossa strenua* (Cresson) (both Apidae) are native to North America and are present in much of the Mid-Atlantic (Hurd et al. 1971). *Peponapis pruinosa* is present throughout the Mid-Atlantic while *X. strenua* is restricted to Virginia and southern Maryland (Hurd et al. 1971). Both species are oligoleges of plants in the genus *Cucurbita* L. (gourds), which only contains adventive species in the Mid-Atlantic. However, pumpkin and squash are extensively cultivated each year and both species are established and common in farms and home gardens. Both species may well have been established in the region for centuries, since squash and gourds were domesticated and widely planted by eastern Native American populations (Price 2009). *Chelostoma philadelphia* (Robertson) (Megachilidae) is the only native *Chelostoma* Latreille species found in eastern North America and has long been associated with the introduced woody shrub *Philadelphus coronarius* L., sweet mock orange. *Philadelphus* L. species are native to the Mid-Atlantic, but such plants are extremely uncommon. Mike Arduser (pers. comm.) and Sam Droege (pers. comm.) have records for *C. philadelphia* in areas where cultivated and wild *Philadelphus* plants do not occur nearby, suggesting unrestricted use of *Philadelphus*. Mellitologists are encouraged to research *C. philadelphia* pollen preferences because several *Chelostoma* species are known pollen specialists elsewhere in the world.

Maryland's bees are relatively well known within the Mid-Atlantic and approximately 21% (92 spp.) of Maryland bee species (431 spp.) are cleptoparasitic (laying eggs in the nests of other bees) (Droege, unpublished data) and cannot be pollen specialists since they gather no pollen. However, some cleptoparasitic bees are associated with specialist bees and therefore are indirectly dependent on the same plant groups as the specialist bees. For example, *Epeoloides pilosula* (Cresson) (Apidae), a nest parasite of now very uncommon *Macropis* species which specialize on *Lysimachia*, is extremely rare and of great conservation concern (Xerces Society 2015). One hundred six of the 131

documented specialist bees occur in Maryland based on extensive unpublished studies of regional bee species distributions (Droege, unpublished data). Such specialist bees comprise 31% of the pollen-carrying species (non-cleptoparasitic) in Maryland (339 spp.). A similar ratio is probably applicable to the entire Mid-Atlantic region.

Plants

Conservation and enhancement of native forbs, particularly *Solidago*, *Symphyotrichum*, *Helianthus*, *Euthamia* Nutt. ex Cass. (goldentops), *Oenothera*, and *Physalis* L. (groundcherries) should be fundamental for the maintenance and restoration of plant-pollinator interactions in fields, meadows, roadsides, and waste areas. The maintenance of diverse *Vaccinium*, *Lyonia* Nutt. (staggerbushes), and *Rhododendron* L. (rhododendrons) shrub communities in acidic soils will benefit many bee species. Woodland host plant restoration is also essential because many specialist bee species only live in rich deciduous forests and forage spring ephemerals (e.g., *Claytonia* L. [springbeauties] and *Erythronium* L. [fawnlilies]) which are absent in fields. Moreover, wetland restoration plantings should include *Hibiscus* L. (rosemallows), *Lysimachia*, *Pontederia* L. (pickerelweeds), *Salix*, and ericaceous plants to provide forage for known and possible wetland oligoleges.

Introduced plants should be avoided or counter-balanced by conservation and propagation of Mid-Atlantic native host plants. Native host plants are often absent from standard planting mixes, which do increase generalist bee abundance, but do not augment specialist bee biodiversity. Highly attractive and productive introduced selections should not be eliminated from plantings, but should perform more supporting roles.

This manuscript serves as a call to regional native plant propagators: add an abundance and richness of specialist bee host plants to nursery stock and seed banks. Although a number of specialist bee host plants are available from retail nurseries and seed distributors, many ecotypes of host plant genera or species remain uncultivated. Accordingly, opportunities are present for Mid-Atlantic plant propagators to diversify plant selections for ecological, social, and economic sustainability.

Future work

Given the disorganized and sparse data on pollen preferences of bees, the difficulty in determining whether bees collected on flowers are provisioning their nests with pollen from that flower or are simply collecting nectar, and that non-provisioning male bees are included in floral records, strict and quantifiable measurements of specialization are currently impossible. Data need to exist for combined floral visitation and pollen consumption records. However, since the purpose of this paper is not a precise quantification of specialization, but a basis for expanding recommended plant lists for pollinator conservation, patterns of specialization, even from disorganized and sparse data, are usually clear.

Some readers may be concerned that compiled records might be incomplete. Any quantitative definition of host plants for bees will be partially incomplete given the broad and imprecise range of available source information on spatiotemporal pollen gathering activities. Available sources almost never provide complete, consistent, or rigorously

quantitative data or details for each bee species. However, we feel that the combination of weighing the literature, information from museum collections, direct field experience, and vetting the catalog with field experts allows us to qualitatively identify these primary associations at the genus level. If this publication stimulates subsequent investigations and clarifications, then this manuscript will be a success.

There is an information gap in our understanding of the status and distribution of bees and associated plant populations in the Mid-Atlantic. Published bee faunas are generally sparse for North America (<50) and primarily of recent origin, particularly in contrast to the >2700 published floras (Palmer 2015). Such a discrepancy between faunas and floras indicates an overall lack of knowledge concerning the taxonomy, life history, and distribution of bees. Eight Maryland, two Pennsylvania, two New Jersey, and one Virginia bee faunas have been published (Winfree et al. 2007 [NJ]; Winfree et al. 2008 [NJ, PA]; Droege et al. 2009 [MD]; Steury et al. 2009 [VA]; Donovall and vanEngelsdorp 2010 [PA]; Orr 2010 [MD]; Droege and Shapiro 2011, 2012 [MD]; Shapiro and Droege 2011, 2012 [MD]; Droege 2013 [MD]; Scarpulla 2013 [MD]). However, only one state, Pennsylvania, has published a comprehensive bee fauna (Donovall and vanEngelsdorp 2010), which is primarily a simple checklist.

Species-level patterns of specialization should be clarified by future collections and research. Degrees and patterns of pollen use by Mid-Atlantic species are undoubtedly more complicated and nuanced than presented here. Listed specialists may forage on alternate pollens at range edges or during resource dearth (e.g., *Habropoda laboriosa* [Fabricius], Southeastern Blueberry Bee, with *Cercis canadensis* L., eastern redbud); specialists might restrict pollen foraging to limited plant species within documented genera; and specialists may be so understudied that some bees likely have failed to be listed and some, perhaps, should be delisted. Consequently, there is still much to be learned about both the distribution of specialist bees in the Mid-Atlantic as well as their pollen gathering preferences.

ACKNOWLEDGMENTS

I thank Samuel W. Droege (United States Geological Survey, Bee Inventory and Monitoring Laboratory), Michael S. Arduser (formerly Missouri Department of Conservation), John S. Ascher (National University of Singapore), Daniel P. Cariveau (University of Minnesota), Charles S. (Charley) Eiseman (freelance naturalist), Jason Gibbs (Michigan State University), Robert P. Jean (Environmental Solutions & Innovations), John L. (Jack) Neff (Central Texas Melittological Institute), T'ai H. Roulston (University of Virginia), and Kimberly A. Stoner (The Connecticut Agricultural Experiment Station) for comments. I greatly appreciate the efforts of the Discover Life and the USDA-NRCS PLANTS Database groups to compile distribution and natural history information for bees and plants respectively.

LITERATURE CITED

Adamson, N.L., B. Borders, J.K. Cruz, S.F. Jordan, K. Gill, J. Hopwood, E. Lee-Ma□der, A. Minnerath, and M. Vaughan. 2015. Pollinator Plants: Mid-Atlantic

- Region. Available at: http://www.xerces.org/wp-content/uploads/2014/09/MidAtlanticPlantList_web.pdf. Accessed 12 March 2016.
- Ascher, J.S., and J. Pickering. 2016. Discover Life bee species guide and world checklist (Hymenoptera: Apoidea: Anthophila). Available at: http://www.discoverlife.org/mp/20q?guide=Apoidea_species. Accessed 15 August 2016.
- Bartomeus, I., J.S. Ascher, J. Gibbs, B.N. Danforth, D.L. Wagner, S.M. Hedtke, and R. Winfree. 2013. Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences of the United States of America* 110(12):4656-4660.
- Bouseman, J.K., and W.E. LaBerge. 1978. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part IX. Subgenus *Melandrena*. *Transactions of the American Entomological Society* 104(3/4):275-389.
- Donovall, L.R., III, and D. vanEngelsdorp. 2010. A checklist of the bees (Hymenoptera: Apoidea) of Pennsylvania. *Journal of the Kansas Entomological Society* 83(1):7-24.
- Droege, S.W. 2013. Investigation of the spring bee fauna (Hymenoptera: Apoidea) of seven woodland sites in the Coastal Plain of Maryland using continuously-trapping arrays of propylene glycol cup traps. *The Maryland Entomologist* 6(1):43-55.
- Droege, S.W., and L.H. Shapiro. 2011. An August survey of wild bees (Hymenoptera: Apoidea) in the northeastern port areas of Baltimore, Maryland and the second North American record of *Pseudoanthidium nanum* (Mocsáry). *The Maryland Entomologist* 5(3):33-44.
- Droege, S.W., and L.H. Shapiro. 2012. An August survey of wild bees (Hymenoptera: Apoidea) in the northeastern port areas of Baltimore, Maryland and the second North American record of *Pseudoanthidium nanum* (Mocsáry) – Addendum. *The Maryland Entomologist* 5(4):30.
- Droege, S., C.A. Davis, W.E. Steiner, Jr., and J. Mawdsley. 2009. The lost micro-deserts of the Patuxent River: Using landscape history, insect and plant specimens, and field work to detect and define a unique community. *Proceedings of the Entomological Society of Washington* 111(1):132-144.
- Cane, J.H., and S. Sipes. 2006. Floral specialization by bees: Analytical methodologies and a revised lexicon for oligolecty. Pages 99-122 in Waser, N.M., and J. Ollerton (Editors), *Plant-Pollinator Interactions: From specialization to generalization*. University of Chicago Press, Chicago, IL. 488 pp.
- Fowler, J. 2016. Specialist bees of the Northeast: host plants and habitat conservation. *Northeastern Naturalist* 23(2):305-320.
- Hilty, J.A. 2012. Oligolectic Bees. Available at: http://www.illinoiswildflowers.info/flower_insects/files/oligoleges.htm. Accessed 1 June 2015.

- Hurd, P.D., Jr., W.E. LaBerge, and E.G. Linsley. 1980. *Principal Sunflower Bees of North America with Emphasis on the Southwestern United States (Hymenoptera: Apoidea)*. Smithsonian Contributions to Zoology Number 310. Smithsonian Institution Press, Washington, DC. 158 pp.
- Hurd, P.D., Jr., E.G. Linsley, and T.W. Whitaker. 1971. Squash and gourd bees (*Peponapis*, *Xenoglossa*) and the origin of the cultivated *Cucurbita*. *Evolution* 25(1):218-234.
- Jones, K.B., K.H. Riitters, J.D. Wickham, R.D. Tankersley, Jr., R.V. O'Neill, D.J. Chaloud, E.R. Smith, and A.C. Neale. 1997. *An Ecological Assessment of the United States Mid-Atlantic Region: A Landscape Atlas*. United States Environmental Protection Agency, Office of Research and Development, Washington DC. EPA/600/R-97/130. 104 pp.
- Krombein, K.V., P.D. Hurd, Jr., D.R. Smith, and B.D. Burks. 1979. *Catalog of Hymenoptera in America North of Mexico*. 3 Volumes. Smithsonian Institution Press, Washington, DC. 2735 pp.
- LaBerge, W.E. 1967. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part I. *Callandrena* (Hymenoptera: Andrenidae). *Bulletin of the University of Nebraska State Museum* 7:1-316.
- LaBerge, W.E. 1969. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part II. *Plastandrena*, *Aporandrena*, *Charitandrena*. *Transactions of the American Entomological Society* 95(1):1-47.
- LaBerge, W.E. 1971. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part IV. *Scapteropsis*, *Xiphandrena* and *Rhaphandrena*. *Transactions of the American Entomological Society* 97(3):441-520.
- La Berge [sic], W.E. 1973. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part VI. Subgenus *Trachandrena*. *Transactions of the American Entomological Society* 99(3):235-371.
- La Berge [sic], W.E. 1977. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part VIII. Subgenera *Thysandrena*, *Dasyandrena*, *Psammandrena*, *Rhacandrena*, *Euandrena*, *Oxyandrena*. *Transactions of the American Entomological Society* 103(1):1-143.
- LaBerge, W.E. 1980. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part X. Subgenus *Andrena*. *Transactions of the American Entomological Society* 106(4):395-525.
- LaBerge, W.E. 1985. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part XI. Minor subgenera and subgeneric key. *Transactions of the American Entomological Society* 111(4):441-567.

- LaBerge, W.E. 1986. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part XII. Subgenera *Leucandrena*, *Ptilandrena*, *Scoliandrena* and *Melandrena*. *Transactions of the American Entomological Society* 112(3):191-248.
- LaBerge, W.E. 1989. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part XIII. Subgenera *Simandrena* and *Taeniandrena*. *Transactions of the American Entomological Society* 115(1):1-56.
- LaBerge, W.E., and J.K. Bouseman. 1970. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part III. *Tylandrena*. *Transactions of the American Entomological Society* 96(4):543-605.
- LaBerge, W.E., and D.W. Ribble. 1972. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part V. *Gonandrena*, *Geissandrena*, *Parandrena*, *Pelicanandrena*. *Transactions of the American Entomological Society* 98(3):271-358.
- LaBerge, W.E., and D.W. Ribble. 1975. A revision of the bees of the genus *Andrena* of the Western Hemisphere. Part VII. Subgenus *Euandrena*. *Transactions of the American Entomological Society* 101(3):371-446.
- Linsley, E.G., and MacSwain, J.W. 1958. The significance of floral constancy among bees of the genus *Diadasia* (Hymenoptera, Anthophoridae). *Evolution* 12(2):219-223.
- Lovell, J.H. 1918. *The Flower and the Bee: Plant life and pollination*. Charles Scribner's Sons, New York, NY. 286 pp.
- Minckley, R.L., W.T. Wcislo, D. Yanega, and S.L. Buchmann. 1994. Behavior and phenology of a specialist bee (*Dieunomia*) and sunflower (*Helianthus*) pollen availability. *Ecology* 75(5):1406-1419.
- Mitchell, T.B. 1960. *Bees of the Eastern United States*, Volume I. The North Carolina Agricultural Experiment Station, Technical Bulletin Number 141, Raleigh, NC. 538 pp.
- Mitchell, T.B. 1962. *Bees of the Eastern United States*, Volume II. The North Carolina Agricultural Experiment Station, Technical Bulletin Number 152, Raleigh, NC. 557 pp.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. Map Supplement (scale 1:7,500,000). *Annals of the Association of American Geographers* 77(1):118-125.
- Orr, R.L. 2010. Preliminary list of the bees (Hymenoptera: Apoidea) of Assateague Island National Seashore, Worcester County, Maryland. *The Maryland Entomologist* 5(2):41-50.
- Palmer, M.W. 2015. The floras of North America project. Available at <http://botany.okstate.edu/floras/>. Accessed 1 June 2015.

- Price, T.D. 2009. Ancient farming in eastern North America. *Proceedings of the National Academy of Sciences of the United States of America* 106(16):6427-6428.
- Radeloff, V.C., R.B. Hammer, S.I. Stewart, J.S. Fried, S.S. Holcomb, and J.F. McKeefry. 2005. The wildland-urban interface in the United States. *Ecological Applications* 15(3):799-805.
- Rafferty, N.E., P.J. CaraDonna, and J.L. Bronstein. 2015. Phenological shifts and the fate of mutualisms. *Oikos*, 124(1):14-21.
- Robertson, C. 1925. Heterotropic bees. *Ecology* 6(4):412-436.
- Robertson, C. 1926. Revised list of oligolectic bees. *Ecology* 7(3):378-380.
- Robertson, C. 1929. Phenology of oligolectic bees and favorite flowers. *Psyche* 36(2):112-118.
- Scarpulla, E.J. 2013. A yearlong survey of the bees (Hymenoptera: Apoidea) of a human-made habitat created from dredged material: Hart-Miller Island, Chesapeake Bay, Baltimore County, Maryland. *The Maryland Entomologist* 6(1):56-84.
- Shapiro, L.H., and S.W. Droege. 2011. Bees (Hymenoptera: Apoidea) of the Dominion Cove Point Liquefied Natural Gas Facility and vicinity, Calvert County, Maryland. *The Maryland Entomologist* 5(3):45-65.
- Shapiro, L.H., and S.W. Droege. 2012. Bees (Hymenoptera: Apoidea) of the Dominion Cove Point Liquefied Natural Gas Facility and vicinity, Calvert County, Maryland – Addendum. *The Maryland Entomologist* 5(4):31.
- Steury, B.W., S.W. Droege, and E.T. Oberg. 2009. Bees (Hymenoptera: Anthophila) of a riverside outcrop prairie in Fairfax County, Virginia. *Banisteria* 34:17-24.
- Winfree, R., N.M. Williams, H. Gaines, J.S. Ascher, and C. Kremen. 2008. Wild bee pollinators provide the majority of crop visitation across land-use gradients in New Jersey and Pennsylvania, USA. *Journal of Applied Ecology* 45(3):793-802.
- Winfree, R., T. Griswold, and C. Kremen. 2007. Effect of human disturbance on bee communities in a forested ecosystem. *Conservation Biology* 21(1):213-223.
- USDA-NRCS (United States Department of Agriculture - Natural Resources Conservation Service). 2016. The PLANTS Database. National Plant Data Team, Greensboro, NC. Available at: <http://plants.usda.gov/java/>. Accessed 06-07 March 2016.
- Xerces Society for Invertebrate Conservation, The. 2015. Red list of bees: Native bees in decline. Available at: <http://www.xerces.org/pollinator-redlist/>. Accessed 1 June 2015.

Table 3 (Supplementary). Mid-Atlantic native specialist bee families (n = 6), genera (n = 25), and species (n = 131). Scientific names (*Genus* [*Subgenus*] *species*, Taxonomic authority), native ranges according to records from the six Mid-Atlantic states (DE, MD, NJ, PA, VA, and WV), seasons of peak activity (Spring, Summer, or Fall), local conservation status (Common = C, Uncommon = U, or Rare = R), and host plant genera or host plant families are provided. Bee families and member genera and species are presented in ascending alphabetical order. Peak activity was provided by Droege (in litt.) and associated with host plant flowering phenology. Status was also provided by Droege (in litt.). All other data were compiled initially from Discover Life (Ascher and Pickering 2016), checked against published sources (e.g., Krombein et al. 1979, and references therein) if available, refined through discussions with regional bee experts, and supplemented and revised with unpublished information provided by John S. Ascher (in litt.).

Family	Scientific Name	Native Range	Peak Activity	Status	Host Plant Genus or Family
Andrenidae	<i>Andrena</i> (<i>Callandrena</i>) <i>accepta</i> Viereck, 1916	MD-NJ-VA	Fall	R	<i>Helianthus</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>aliciae</i> Robertson, 1891	DE-MD-NJ-PA-VA-WV	Summer	R	<i>Helianthus</i>
	<i>Andrena</i> (<i>Parandrena</i>) <i>andrenoides</i> (Cresson, 1878)	MD-NJ-VA	Spring	R	<i>Salix</i>
	<i>Andrena</i> (<i>Scaphandrena</i>) <i>arabis</i> Robertson, 1897	DE-MD-NJ-PA-VA-WV	Spring	U	<i>Arabis</i> , <i>Cardamine</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>asteris</i> Robertson, 1891	DE-MD-NJ-PA-VA-WV	Fall	C	<i>Solidago</i> , <i>Symphytotrichum</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>asteroides</i> Mitchell, 1960	MD-NJ-PA-VA	Fall	R	<i>Symphytotrichum</i>
	<i>Andrena</i> (<i>Thysandrena</i>) <i>bisalicis</i> Viereck, 1908	DE-MD-NJ-PA-VA-WV	Spring	U	<i>Salix</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>braccata</i> Viereck, 1907	MD-NJ-PA-VA	Fall	R	<i>Euthamia</i> , <i>Solidago</i>
	<i>Andrena</i> (<i>Conandrena</i>) <i>bradleyi</i> Viereck, 1907	DE-MD-NJ-PA-VA	Spring	U	<i>Gaylussacia</i> , <i>Vaccinium</i>
	<i>Andrena</i> (<i>Cnemidandrena</i>) <i>canadensis</i> Dalla Torre, 1896	NJ-VA	Fall	U	<i>Solidago</i>
	<i>Andrena</i> (<i>Andrena</i>) <i>carolina</i> Viereck, 1909	DE-MD-NJ-PA-VA-WV	Spring	C	<i>Vaccinium</i>
	<i>Andrena</i> (<i>Cnemidandrena</i>) <i>chromotricha</i> Cockerell, 1899	PA-WV	Fall	R	<i>Asteraceae</i>
	<i>Andrena</i> (<i>Andrena</i>) <i>clarkella</i> (Kirby, 1802)	WV	Spring	R	<i>Salix</i>
	<i>Andrena</i> (<i>Andrena</i>) <i>cornelli</i> Viereck, 1907	MD-NJ-PA-VA	Spring	U	<i>Rhododendron</i>
	<i>Andrena</i> (<i>Ptilandrena</i>) <i>distans</i> Provancher, 1888	MD-NJ-PA-VA	Spring	U	<i>Geranium</i>

Family	Scientific Name	Native Range	Peak Activity	Status	Host Plant Genus or Family
Andrena	(<i>Callandrena</i>) <i>duplicata</i> Mitchell, 1960	NJ	Fall	R	Asteraceae
	<i>Andrena</i> (<i>Ptilandrena</i>) <i>erigeniae</i> Robertson, 1891	DE-MD-NJ-PA-VA-WV	Spring	C	<i>Claytonia</i>
	<i>Andrena</i> (<i>Tylandrena</i>) <i>erythrogaster</i> (Ashmead, 1890)	MD-NJ-PA-VA-WV	Spring	U	<i>Salix</i>
	<i>Andrena</i> (<i>Leucandrena</i>) <i>erythronii</i> Robertson, 1891	MD-NJ-PA-VA-WV	Spring	U	<i>Erythronium</i>
	<i>Andrena</i> (<i>Gonandrena</i>) <i>fragilis</i> Smith, 1853	DE-MD-NJ-PA-VA-WV	Summer	C	<i>Cornus</i>
	<i>Andrena</i> (<i>Andrena</i>) <i>frigida</i> Smith 1853	MD-NJ-PA-VA	Spring	U	<i>Salix</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>fulvipennis</i> Smith, 1853	DE-MD-NJ	Fall	R	<i>Chrysopsis</i> , <i>Heterotheca</i> , <i>Pityopsis</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>gardineri</i> Cockerell, 1906	MD-VA-WV	Spring	U	<i>Packera</i>
	<i>Andrena</i> (<i>Euandrena</i>) <i>geranii</i> , Robertson, 1891	MD-NJ-PA-VA-WV	Spring	U	<i>Hydrophyllum</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>helianthi</i> Robertson, 1891	NJ-PA-VA	Summer	R	<i>Helianthus</i>
	<i>Andrena</i> (<i>Chemidandrena</i>) <i>hirticincta</i> Provancher, 1888	MD-NJ-PA-VA-WV	Fall	C	<i>Euthamia</i> , <i>Solidago</i>
	<i>Andrena</i> (<i>Micrandrena</i>) <i>illinoensis</i> Robertson, 1891	MD-VA	Spring	R	<i>Salix</i>
	<i>Andrena</i> (<i>Gonandrena</i>) <i>integra</i> Smith, 1853	MD-NJ-PA-VA	Summer	U	<i>Cornus</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>krigiana</i> Robertson, 1901	MD-NJ-PA-VA-WV	Spring	U	<i>Krigia</i>
	<i>Andrena</i> (<i>Micrandrena</i>) <i>lamelliterga</i> Ribble, 1968	MD-PA-VA	Spring	R	<i>Phacelia</i>
	<i>Andrena</i> (<i>Andrena</i>) <i>macoupinensis</i> Robertson, 1900	MD-PA-VA	Spring	U	<i>Salix</i>
	<i>Andrena</i> (<i>Trachandrena</i>) <i>mariae</i> Robertson, 1891	MD-NJ-VA-WV	Spring	U	<i>Salix</i>
	<i>Andrena</i> (<i>Micrandrena</i>) <i>melanochroa</i> Cockerell, 1898	MD-NJ-VA-WV	Spring	U	Rosaceae, e.g., <i>Fragaria</i>
	<i>Andrena</i> (<i>Parandrena</i>) <i>nida</i> Mitchell, 1960	MD-NJ-VA-WV	Spring	U	<i>Salix</i>
	<i>Andrena</i> (<i>Micrandrena</i>) <i>nigrae</i> Robertson, 1905	DE-MD-NJ-PA-VA-WV	Spring	U	<i>Salix</i>
Andrena	(<i>Chemidandrena</i>) <i>nubecula</i> Smith, 1853	DE-MD-NJ-PA-VA-WV	Fall	U	<i>Solidago</i> , <i>Symphytotrichum</i>
	<i>Andrena</i> (<i>Euandrena</i>) <i>phaceliae</i> Mitchell, 1960	MD-VA-WV	Spring	R	<i>Phacelia</i>
	<i>Andrena</i> (<i>Callandrena</i> s.l.) <i>placata</i> Mitchell, 1960	MD-NJ-PA-VA-WV	Fall	C	<i>Solidago</i> , <i>Symphytotrichum</i>
	<i>Andrena</i> (<i>Gonandrena</i>) <i>platyparia</i> Robertson, 1895	MD-NJ-PA-VA	Summer	U	<i>Cornus</i>

Family	Scientific Name	Native Range	Peak Activity	Status	Host Plant Genus or Family
	<i>Andrena (Euandrena) polemonii</i> Robertson, 1891	WV	Spring	R	<i>Polemonium</i>
	<i>Andrena (Callandrena s.l.) rudbeckiae</i> Robertson, 1891	DE-MD-NJ-PA	Summer	R	<i>Rudbeckia</i>
	<i>Andrena (Micrandrena) salicaria</i> Robertson, 1905	MD-NJ-PA-VA-WV	Spring	R	<i>Salix</i>
	<i>Andrena (Callandrena s.l.) simplex</i> Smith, 1853	DE-MD-NJ-PA-VA-WV	Fall	C	<i>Solidago, Symphyotrichum</i>
	<i>Andrena (Derandrena) ivulariae</i> Mitchell, 1960	MD-NJ-PA-VA-WV	Spring	R	<i>Uvularia</i>
	<i>Andrena (Iomelissa) violae</i> Robertson, 1891	DE-MD-NJ-PA-VA-WV	Spring	C	<i>Viola</i>
	<i>Andrena (Micrandrena) ziziae</i> Robertson, 1891	MD-NJ-PA-VA-WV	Spring	U	<i>Zizia</i>
	<i>Andrena (Derandrena) ziziaeformis</i> Cockerell, 1908	DE-MD-NJ-PA-VA-WV	Spring	U	<i>Potentilla, Waldsteinia</i>
	<i>Calliopsis (Verbenapis) nebraskensis</i> Crawford, 1902	NJ	Summer	R	<i>Verbena</i>
	<i>Panurginus atramentosus</i> Crawford, 1926	MD-NJ-VA	Spring	R	<i>Vaccinium</i>
	<i>Panurginus potentillae</i> (Crawford, 1916)	MD-NJ-VA	Spring	U	<i>Potentilla</i>
	<i>Perdita (Cockerellia) bequaerti</i> Viereck, 1917	MD-NJ	Summer	R	Asteraceae
	<i>Perdita (Hexaperdita) bishoppi</i> Cockerell, 1906	DE-MD-NJ	Fall	R	<i>Chrysopsis, Heterotheca, Pityopsis</i>
	<i>Perdita (Hexaperdita) boltoniae</i> (Robertson, 1902)	DE-MD-NJ-VA	Fall	R	<i>Chrysopsis, Heterotheca, Pityopsis</i>
	<i>Perdita (Perdita) consobrina</i> Timberlake, 1928	VA	Fall	R	Asteraceae
	<i>Perdita (Perdita) gerardiae</i> Crawford, 1932	MD-NJ	Fall	R	<i>Agalinis</i>
	<i>Perdita (Perdita) gerhardi</i> Viereck, 1904	MD-NJ-VA-WV	Summer	R	<i>Monarda</i>
	<i>Perdita (Perdita) halictoides</i> Smith, 1853	MD-NJ	Summer	R	<i>Physalis</i>
	<i>Perdita (Perdita) octomaculata</i> (Say, 1824)	DE-MD-NJ-VA-WV	Fall	U	<i>Solidago</i>
	<i>Perdita (Perdita) swenki</i> Crawford, 1915	NJ	Summer	R	Asteraceae
	<i>Protandrena abdominalis</i> (Cresson, 1878)	MD-NJ-PA-VA-WV	Summer	R	<i>Monarda</i>
	<i>Pseudopanurgus aestivalis</i> (Provancher, 1882)	MD	Fall	R	<i>Solidago, Symphyotrichum</i>
	<i>Pseudopanurgus alb tarsis</i> (Cresson, 1872)	VA	Summer	R	Asteraceae
	<i>Pseudopanurgus andrenoides</i> (Smith, 1853)	NJ-VA-WV	Fall	U	Asteraceae

Family	Scientific Name	Native Range	Peak Activity	Status	Host Plant Genus or Family
Apidae	<i>Pseudopanurgus compositarum</i> (Robertson, 1893)	DE-MD-NJ-PA-VA-WV	Fall	U	Asteraceae
	<i>Pseudopanurgus illinoiensis</i> (Cresson, 1878)	MD-VA	Spring	R	Asteraceae
	<i>Pseudopanurgus labrosiformis</i> (Robertson, 1898)	MD-WV	Summer	R	Asteraceae
	<i>Pseudopanurgus labrosus</i> (Robertson, 1895)	VA-WV	Summer	U	Asteraceae
	<i>Pseudopanurgus pauper</i> (Cresson, 1878)	MD-NJ-PA-VA-WV	Summer	R	<i>Ceanothus</i>
	<i>Pseudopanurgus rugosus</i> (Robertson, 1895)	MD-VA	Summer	U	Asteraceae
	<i>Pseudopanurgus virginicus</i> (Cockerell, 1907)	MD-VA-WV	Spring	R	<i>Houstonia</i>
	<i>Anthophorula (Anthophorisca) micheneri</i> Timberlake, 1948	VA	Fall	R	<i>Agalinis</i>
	<i>Cemolobus ipomoeae</i> (Robertson, 1891)	MD-PA-VA-WV	Summer	R	<i>Ipomoea</i>
	<i>Florilegus (Florilegus) condignus</i> (Cresson, 1878)	MD-NJ-VA	Summer	R	<i>Pontederia</i>
	<i>Habropoda laboriosa</i> (Fabricius, 1804)	DE-MD-NJ-PA-VA	Spring	C	<i>Cercis, Vaccinium</i>
	<i>Melissodes (Eumelissodes) agilis</i> Cresson, 1878	DE-MD-NJ-PA-VA-WV	Summer	C	Asteraceae, e.g., <i>Helianthus</i>
	<i>Melissodes (Apomelissodes) apicatus</i> Lovell & Cockerell, 1906	DE-MD-NJ	Summer	U	<i>Pontederia</i>
	<i>Melissodes (Eumelissodes) boltoniae</i> Robertson, 1905	DE-MD-PA-VA	Fall	U	Asteraceae
	<i>Melissodes (Eumelissodes) denticulatus</i> Smith, 1854	DE-MD-NJ-PA-VA-WV	Fall	U	Asteraceae, e.g., <i>Vernonia</i>
	<i>Melissodes (Eumelissodes) dentiventris</i> Smith, 1854	DE-MD-NJ-PA-VA	Fall	U	Asteraceae
	<i>Melissodes (Heliomelissodes) desponsus</i> Smith, 1854	DE-MD-NJ-PA-VA-WV	Fall	C	Asteraceae, e.g., <i>Cirsium</i>
	<i>Melissodes (Eumelissodes) druriellus</i> (Kirby, 1802)	MD-NJ-PA-VA-WV	Fall	U	Asteraceae
	<i>Melissodes (Apomelissodes) fimbriatus</i> Cresson, 1878	VA	Summer	R	<i>Oenothera</i>
	<i>Melissodes (Eumelissodes) fumosus</i> LaBerge, 1961	MD-VA	Fall	R	Asteraceae
	<i>Melissodes (Eumelissodes) illatus</i> Lovell & Cockerell, 1906	WV	Fall	U	Asteraceae
	<i>Melissodes (Eumelissodes) niveus</i> Robertson, 1895	MD-NJ-VA	Fall	U	Asteraceae
	<i>Melissodes (Eumelissodes) subillatus</i> LaBerge, 1961	DE-MD-NJ-PA-VA-WV	Fall	U	Asteraceae
	<i>Melissodes (Eumelissodes) trinodis</i> Robertson, 1901	DE-MD-NJ-VA	Fall	U	Asteraceae

Family	Scientific Name	Native Range	Peak Activity	Status	Host Plant Genus or Family
Colletidae	<i>Melittoma taurea</i> (Say, 1837)	DE-MD-NJ-PA-V-A-WV	Summer	C	<i>Ipomoea</i>
	<i>Ptilothrix bombiformis</i> (Cresson, 1878)	DE-MD-NJ-PA-V-A-WV	Summer	C	<i>Hibiscus</i>
	<i>Svastra (Anthedonia) compta</i> (Cresson, 1878)	MD-NJ-PA	Summer	U	<i>Oenothera</i>
	<i>Svastra (Epimelissodes) obliqua</i> (Say, 1837)	DE-MD-NJ-V-A	Summer	U	Asteraceae
	<i>Svastra (Epimelissodes) petulca</i> (Cresson, 1878)	NJ	Summer	R	Asteraceae
	<i>Colletes aestivalis</i> Patton, 1879	DE-MD-V-A	Summer	R	<i>Heuchera</i>
	<i>Colletes banksi</i> Swenk, 1908	NJ-V-A	Summer	R	<i>Ilex</i>
	<i>Colletes brevicornis</i> Robertson, 1897	MD-NJ-V-A-WV	Spring	U	<i>Campanula, Triodanis</i>
	<i>Colletes compactus</i> Cresson, 1868	DE-MD-NJ-PA-V-A-WV	Fall	C	Asteraceae
	<i>Colletes latitarsis</i> Robertson, 1891	MD-NJ-PA-V-A-WV	Summer	U	<i>Physalis</i>
	<i>Colletes productus</i> Robertson, 1891	MD-NJ-V-A-WV	Summer	R	<i>Lyonia, Vaccinium</i>
	<i>Colletes simulans</i> Cresson, 1868	MD-NJ-PA-V-A-WV	Fall	C	<i>Euthamia, Solidago, Symphyotrichum</i>
	<i>Colletes solidaginis</i> Swenk, 1906	MD-NJ-V-A	Fall	R	<i>Solidago</i>
	<i>Colletes speculiferus</i> Cockerell, 1927	DE-MD-NJ-V-A	Fall	U	Asteraceae
	<i>Colletes thysanellae</i> Mitchell, 1951	V-A	Fall	R	Asteraceae
Halictidae	<i>Colletes validus</i> Cresson, 1868	DE-MD-NJ-PA-V-A	Spring	U	Ericaceae
	<i>Colletes willistoni</i> Robertson, 1891	MD-NJ-PA-V-A	Summer	R	<i>Physalis</i>
	<i>Dieunomia (Dieunomia) heteropoda</i> (Say, 1824)	MD-NJ-V-A	Fall	R	<i>Bidens, Helianthus</i>
	<i>Dieunomia (Epinomia) nevadensis</i> (Cresson, 1874)	MD	Summer	R	Asteraceae
	<i>Dufourea monardae</i> (Viereck, 1924)	NJ	Summer	R	<i>Monarda</i>
	<i>Dufourea novaeangliae</i> (Robertson, 1897)	MD-NJ-PA-V-A	Summer	R	<i>Pontederia</i>
	<i>Lasioglossum (Hemihalictus) lustrans</i> (Cockerell, 1897)	DE-MD-V-A	Summer	R	<i>Pyrrhopappus</i>
	<i>Lasioglossum (Sphecodogastra) oenotherae</i> (Stevens, 1920)	MD-NJ-PA-V-A-WV	Summer	U	<i>Oenothera</i>
	<i>Lasioglossum (Hemihalictus) pectinatum</i> (Robertson, 1890)	DE-MD-NJ-PA-V-A-WV	Summer	R	<i>Physalis</i>

Family	Scientific Name	Native Range	Peak Activity	Status	Host Plant Genus or Family
Megachilidae	<i>Hoplitis (Robertsonella) simplex</i> (Cresson, 1864)	DE-MD-NJ-VA-WV	Spring	R	<i>Nemophila, Phacelia</i>
	<i>Megachile (Sayapis) inimica</i> Cresson, 1872	DE-MD-NJ-PA-VA-WV	Summer	U	Asteraceae
	<i>Megachile (Megachiloides) integra</i> Cresson, 1878	MD-NJ	Summer	R	<i>Galactia, Strophostyles</i>
	<i>Megachile (Megachiloides) oenotherae</i> (Mitchell, 1924)	NJ	Spring	R	<i>Oenothera</i>
	<i>Megachile (Argyropile) parallela</i> Smith, 1853	VA	Summer	R	Asteraceae
	<i>Megachile (Sayapis) pugnata</i> Say, 1837	DE-MD-NJ-PA-VA-WV	Summer	U	Asteraceae, e.g., <i>Helianthus</i>
	<i>Megachile (Melanosarus) xylocopoides</i> Smith, 1853	DE-MD-NJ-VA	Summer	U	Asteraceae
	<i>Osmia (Helicosmia) chalybea</i> Smith, 1853	DE-MD-NJ-VA-WV	Summer	R	<i>Cirsium</i>
	<i>Osmia (Melanosmia) distincta</i> Cresson, 1864	MD-NJ-PA-VA-WV	Spring	U	<i>Penstemon</i>
	<i>Osmia (Helicosmia) texana</i> Cresson, 1872	MD-VA-WV	Summer	R	<i>Cirsium</i>
	<i>Osmia (Melanosmia) virga</i> Sandhouse, 1939	DE-MD-NJ-PA-VA-WV	Spring	U	<i>Vaccinium</i>
	<i>Paranthidium (Paranthidium) jugatorium</i> (Say, 1824)	MD-NJ-VA-WV	Summer	R	<i>Helianthus</i>
	<i>Trachusa (Heteranthidium) dorsalis</i> (Lepeletier, 1841)	NJ	Summer	R	<i>Strophostyles</i>
	<i>Macropis (Macropis) ciliata</i> Patton, 1880	MD-NJ-PA-VA	Summer	R	<i>Lysimachia</i>
	<i>Macropis (Macropis) nuda</i> (Provancher, 1882)	NJ-PA-WV	Summer	R	<i>Lysimachia</i>
Melittidae	<i>Macropis (Macropis) patellata</i> Patton, 1880	MD-NJ-PA-VA	Summer	R	<i>Lysimachia</i>
	<i>Macropis (Macropis) steironematis</i> Robertson, 1891	VA	Summer	R	<i>Lysimachia</i>
	<i>Melitta (Cilissa) americana</i> (Smith, 1853)	NJ	Summer	R	<i>Vaccinium</i>
	<i>Melitta (Cilissa) eickworthi</i> Snelling & Stage, 1995	MD-NJ-WV	Spring	R	<i>Vaccinium</i>
	<i>Melitta (Cilissa) melittoides</i> (Viereck, 1909)	MD-NJ-PA-VA	Summer	R	<i>Lyonia</i>

Table 4 (Supplementary). Mid-Atlantic native specialist bee primary host plant families (n = 28), genera (n = 54), and species (n = 396). Taxonomic authorities, common names, plant groups (D = Herbaceous, flowering dicot, M = Herbaceous, flowering monocot, S = Broad-leaf woody shrub, ST = Broad-leaf woody shrub-tree, and T = Broad-leaf woody tree), habitats (O = Obligate wetland, FW = Facultative wetland, F = Facultative, FU = Facultative Upland, U = Upland, and ‘ ’ = No data), and native ranges according to six states (DE, MD, NJ, PA, VA, and WV) are provided. Plant families and member genera and species are presented in ascending alphabetical order. All information was compiled from the United States Department of Agriculture - Natural Resources Conservation Service PLANTS Database (2016). Table 4 is presented as a baseline for future research, but should be considered incomplete due to the generally adventitious nature of collections and observations of bees on plants. Bee species attraction to individual plant species within any genus likely varies based on resource availability, quality, and quantity.

Family	Scientific Name	Common Name	Group	Habitat	Range
Apiaceae	<i>Zizia aptera</i> (A. Gray) Fernald	Meadow zizia	D	F	DE-MD-NJ-PA-VA-WV
	<i>Zizia aurea</i> (L.) W.D.J. Koch	Golden zizia	D	F	DE-MD-NJ-PA-VA-WV
	<i>Zizia trifoliata</i> (Michx.) Fernald	Meadow alexanders	D	F-FU	MD-VA-WV
	<i>Ilex ambigua</i> (Michx.) Torr.	Carolina holly	ST	.	VA
Aquifoliaceae	<i>Ilex collina</i> Alexander	Longstalk holly	ST	FW	VA-WV
	<i>Ilex coriacea</i> (Pursh) Chapm.	Large gallberry	ST	FW	VA
	<i>Ilex decidua</i> Walter	Possumhaw	ST	FW	MD-VA
	<i>Ilex glabra</i> (L.) A. Gray	Inkberry	S	FW-F	DE-MD-NJ-PA-VA
	<i>Ilex laevigata</i> (Pursh) A. Gray	Smooth winterberry	ST	O-FW	DE-MD-NJ-PA-VA
	<i>Ilex longipes</i> Chapm. ex Trel.	Georgia holly	ST	F	VA
	<i>Ilex montana</i> Torr. & A. Gray ex A. Gray	Mountain holly	ST	F-FU	MD-NJ-PA-VA-WV
	<i>Ilex mucronata</i> (L.) Powell, Savolainen & Andrews	Catberry	ST	O	MD-NJ-PA-WV
	<i>Ilex opaca</i> Aiton	American holly	ST	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Ilex verticillata</i> (L.) A. Gray	Common winterberry	ST	FW	DE-MD-NJ-PA-VA-WV
	<i>Ilex vomitoria</i> Aiton	Yaupon	ST	F	VA

Family	Scientific Name	Common Name	Group	Habitat	Range
Asteraceae	<i>Bidens alba</i> (L.) DC.	Romerillo	D	.	PA
	<i>Bidens aristosa</i> (Michx.) Britton	Bearded beggarticks	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Bidens beckii</i> Torr. ex Spreng.	Beck's water-marigold	D	O	MD-NJ-PA
	<i>Bidens bidentoides</i> (Nutt.) Britton	Delmarva beggarticks	D	FW	DE-MD-NJ-PA
	<i>Bidens bipinnata</i> L.	Spanish needles	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Bidens cernua</i> L.	Nodding beggartick	D	O	DE-MD-NJ-PA-VA-WV
	<i>Bidens connata</i> Muhl. ex Willd.	Purplestem beggarticks	D	.	DE-MD-NJ-PA-VA-WV
	<i>Bidens coronata</i> (L.) Britton	Crowned beggarticks	D	O	DE-MD-NJ-PA-VA-WV
	<i>Bidens discoides</i> (Torr. & A. Gray) Britton	Small beggarticks	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Bidens eatonii</i> Fernald	Eaton's beggarticks	D	FW	NJ
	<i>Bidens frondosa</i> L.	Devil's beggartick	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Bidens laevis</i> (L.) Britton, Stems & Poggenb.	Smooth beggartick	D	O	DE-MD-NJ-PA-VA-WV
	<i>Bidens mitis</i> (Michx.) Sherff	Smallfruit beggarticks	D	O	DE-MD-NJ-VA
	<i>Bidens tripartita</i> L.	Three-lobed beggarticks	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Bidens vulgata</i> Greene	Big devils beggartick	D	O-F	DE-MD-NJ-PA-VA-WV
	<i>Chrysopsis gossypina</i> (Michx.) Elliott	Cottony goldenaster	D	FU-U	VA
	<i>Chrysopsis mariana</i> (L.) Elliott	Maryland goldenaster	D	U	DE-MD-NJ-PA-VA-WV
	<i>Chrysopsis pilosa</i> Nutt.	Soft goldenaster	D	.	VA
	<i>Cirsium altissimum</i> (L.) Hill	Tall thistle	D	.	DE-MD-NJ-PA-VA-WV
	<i>Cirsium carolinianum</i> (Walter) Fernald & B.G. Schub.	Soft thistle	D	.	VA
Cirsium	<i>Cirsium discolor</i> (Muhl. ex Willd.) Spreng.	Field thistle	D	U	DE-MD-NJ-PA-VA-WV
	<i>Cirsium horridulum</i> Michx.	Yellow thistle	D	F-FU	DE-MD-NJ-PA-VA
	<i>Cirsium muticum</i> Michx.	Swamp thistle	D	O	DE-MD-NJ-PA-VA-WV
	<i>Cirsium nuttallii</i> DC.	Nuttall's thistle	D	F	VA
	<i>Cirsium pumilum</i> Spreng.	Pasture thistle	D	.	DE-MD-NJ-PA-VA-WV
	<i>Cirsium undulatum</i> (Nutt.) Spreng.	Wavyleaf thistle	D	F-FU	PA

Family	Scientific Name	Common Name	Group	Habitat	Range
	<i>Cirsium virginianum</i> (L.) Michx.	Virginia thistle	D	FW	DE-NJ-VA
	<i>Euthamia caroliniana</i> (L.) Greene ex Porter & Britton	Slender goldentop	D	F	DE-MD-NJ-PA-VA
	<i>Euthamia graminifolia</i> (L.) Nutt.	Flat-top goldentop	D	F	DE-MD-NJ-PA-VA-VA-WV
	<i>Euthamia gymnospermoides</i> Greene	Texas goldentop	D	F	WV
	<i>Helianthus angustifolius</i> L.	Swamp sunflower	D	FW	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus annuus</i> L.	Common sunflower	D	F	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus atrorubens</i> L.	Purpldisk sunflower	D	.	NJ-VA
	<i>Helianthus debilis</i> Nutt.	Cucumberleaf sunflower	D	F-U	MD-PA-VA-WV
	<i>Helianthus decapetalus</i> L.	Thinleaf sunflower	D	FU-U	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus divaricatus</i> L.	Woodland sunflower	D	.	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus giganteus</i> L.	Giant sunflower	D	FW	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus grosseserratus</i> M. Martens	Sawtooth sunflower	D	FW-F	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus hirsutus</i> Raf.	Hairy sunflower	D	.	MD-PA-VA-WV
	<i>Helianthus laevigatus</i> Torr. & A. Gray	Smooth sunflower	D	.	MD-PA-VA-WV
	<i>Helianthus maximiliani</i> Schrad.	Maximilian sunflower	D	U	MD-NJ-PA-VA-WV
	<i>Helianthus microcephalus</i> Torr. & A. Gray	Small woodland sunflower	D	U	MD-NJ-PA-VA-WV
	<i>Helianthus mollis</i> Lam.	Ashy sunflower	D	.	MD-NJ-PA-VA-WV
	<i>Helianthus occidentalis</i> Riddell	Fewleaf sunflower	D	FU-U	MD-NJ-PA-VA-WV
	<i>Helianthus pauciflorus</i> Nutt.	Stiff sunflower	D	.	MD-NJ-PA-VA
	<i>Helianthus petiolaris</i> Nutt.	Prairie sunflower	D	.	NJ-PA-VA-WV
	<i>Helianthus strumosus</i> L.	Paleleaf woodland sunflower	D	FU-U	DE-MD-NJ-PA-VA-VA-WV
	<i>Helianthus tuberosus</i> L.	Jerusalem artichoke	D	FU	DE-MD-NJ-PA-VA-VA-WV
	<i>Heterotheca camporum</i> (Greene) Shimmers	Lemonyellow false goldenaster	D	.	NJ-VA
	<i>Heterotheca subaxillaris</i> (Lam.) Britton & Rusby	Camphorweed	D	.	DE-MD-NJ-PA-VA
	<i>Krigia biflora</i> (Walter) S.F. Blake	Twoflower dwarfdandelion	D	FU	DE-MD-NJ-PA-VA-VA-WV
	<i>Krigia caespitosa</i> (Raf.) K.L. Chambers	Weedy dwarfdandelion	D	F	VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
	<i>Krigia dandelion</i> (L.) Nutt.	Potato dwarf dandelion	D	F-FU	MD-NJ-VA
	<i>Krigia virginica</i> (L.) Willd.	Virginia dwarf dandelion	D	FU-U	DE-MD-NJ-PA-VA-WV
	<i>Packera anomia</i> (Alph. Wood) W.A. Weber & Á. Löve	Small's ragwort	D	FU-U	DE-MD-NJ-PA-VA-WV
	<i>Packera aurea</i> (L.) Á. Löve & D. Löve	Golden ragwort	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Packera millefolium</i> (Torr. & A. Gray) W.A. Weber & Á. Löve	Piedmont ragwort	D	.	VA
	<i>Packera obovata</i> (Muhl. ex Willd.) W.A. Weber & Á. Löve	Roundleaf ragwort	D	FU	MD-NJ-PA-VA-WV
	<i>Packera paupercula</i> (Michx.) Á. Löve & D. Löve	Balsam groundsel	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Packera plattensis</i> (Nutt.) W.A. Weber & Á. Löve	Prairie groundsel	D	FU	PA-VA-WV
	<i>Packera tomentosa</i> (Michx.) C. Jeffrey	Woolly ragwort	D	FU	DE-MD-NJ-VA
	<i>Pityopsis aspera</i> (Shuttlw. ex Small) Small	Pineland silkgrass	D	.	VA
	<i>Pityopsis falcata</i> (Pursh) Nutt.	Sickleleaf silkgrass	D	.	NJ
	<i>Pityopsis graminifolia</i> (Michx.) Nutt.	Narrowleaf silkgrass	D	U	DE-MD-VA-WV
	<i>Pyrrhopappus carolinianus</i> (Walter) DC.	Carolina desert-chicory	D	.	DE-MD-PA-VA-WV
	<i>Rudbeckia fulgida</i> Aiton	Orange coneflower	D	F	DE-MD-NJ-PA-VA-WV
	<i>Rudbeckia heliopsisidis</i> Torr. & A. Gray	Sunfacing coneflower	D	.	VA
	<i>Rudbeckia hirta</i> L.	Blackeyed Susan	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Rudbeckia laciniata</i> L.	Cutleaf coneflower	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Rudbeckia triloba</i> L.	Browneyed Susan	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Solidago altissima</i> L.	Canada goldenrod	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Solidago arguta</i> Aiton	Atlantic goldenrod	D	U	DE-MD-NJ-PA-VA-WV
	<i>Solidago bicolor</i> L.	White goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Solidago caesia</i> L.	Wreath goldenrod	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Solidago canadensis</i> L.	Canada goldenrod	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Solidago curtisii</i> Torr. & A. Gray	Mountain decumbent goldenrod	D	FU	MD-PA-VA-WV
	<i>Solidago erecta</i> Pursh	Showy goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Solidago faucibus</i> Wieboldt	Gorge goldenrod	D	.	VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
	<i>Solidago fistulosa</i> Mill.	Pine barren goldenrod	D	FW-F	DE-MD-NJ-VA
	<i>Solidago flaccidifolia</i> Small	Mountain goldenrod	D	.	VA-WV
	<i>Solidago flexicaulis</i> L.	Zigzag goldenrod	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Solidago gigantea</i> Aiton	Giant goldenrod	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Solidago gracillima</i> Torr. & A. Gray	Virginia goldenrod	D	O	VA-WV
	<i>Solidago hispida</i> Muhl. ex Willd.	Hairy goldenrod	D	.	MD-NJ-PA-VA-WV
	<i>Solidago juncea</i> Aiton	Early goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Solidago lanceifolia</i> Torr. & A. Gray	Lance-leaf goldenrod	D	.	VA
	<i>Solidago latissimifolia</i> Mill.	Elliott's goldenrod	D	O	DE-MD-NJ-VA
	<i>Solidago nemoralis</i> Aiton	Gray goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Solidago odora</i> Aiton	Aniscented goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Solidago patula</i> Muhl. ex Willd.	Roundleaf goldenrod	D	O	DE-MD-NJ-PA-VA-WV
	<i>Solidago pinetorum</i> Small	Small's goldenrod	D	.	VA
	<i>Solidago puberula</i> Nutt.	Downy goldenrod	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Solidago radula</i> Nutt.	Western rough goldenrod	D	.	VA
	<i>Solidago roanensis</i> Porter	Roan mountain goldenrod	D	.	MD-NJ-PA-VA-WV
	<i>Solidago rugosa</i> Mill.	Wrinkleleaf goldenrod	D	F	DE-MD-NJ-PA-VA-WV
	<i>Solidago rupestris</i> Raf.	Rock goldenrod	D	.	MD-PA-VA
	<i>Solidago sempervirens</i> L.	Seaside goldenrod	D	FW	DE-MD-NJ-PA-VA
	<i>Solidago simplex</i> Kunth	Mt. Albert goldenrod	D	FU	MD-PA-VA-WV
	<i>Solidago speciosa</i> Nutt.	Showy goldenrod	D	.	MD-NJ-PA-VA-WV
	<i>Solidago sphacelata</i> Raf.	Autumn goldenrod	D	.	VA-WV
	<i>Solidago squarrosa</i> Nutt.	Stout goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Solidago stricta</i> Aiton	Wand goldenrod	D	O-FW	DE-MD-NJ-VA
	<i>Solidago tortifolia</i> Elliott	Twistleaf goldenrod	D	.	VA
	<i>Solidago uliginosa</i> Nutt.	Bog goldenrod	D	O	DE-MD-NJ-PA-VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
	<i>Solidago ulmifolia</i> Muhl. ex Willd.	Elmleaf goldenrod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum boreale</i> (Torr. & Gray) Á. Löve & D. Löve	Northern bog aster	D	O	NJ-PA-WV
	<i>Symphytotrichum ciliatum</i> (Ledeb.) G.L. Nesom	Rayless alkali aster	D	F	PA
	<i>Symphytotrichum concolor</i> (L.) G.L. Nesom	Eastern silver aster	D	.	DE-MD-NJ-VA
	<i>Symphytotrichum cordifolium</i> (L.) G.L. Nesom	Common blue wood aster	D	.	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum depauperatum</i> (Fernald) G.L. Nesom	Serpentine aster	D	.	MD-PA-WV
	<i>Symphytotrichum depauperatum</i> (Fernald) G.L. Nesom	Southern annual saltmarsh aster	D	O	VA
	<i>Symphytotrichum divaricatum</i> (Nutt.) G.L. Nesom	Drummond's aster	D	.	MD-PA-WV
	<i>Symphytotrichum drummondii</i> (Lindl.) G.L. Nesom	Rice button aster	D	F	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum dumosum</i> (L.) G.L. Nesom	March American-aster	D	O	VA
	<i>Symphytotrichum elliotii</i> (Torr. & A. Gray) G.L. Nesom	White heath aster	D	FU-U	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum ericoides</i> (L.) G.L. Nesom	Largeflower aster	D	.	VA
	<i>Symphytotrichum grandiflorum</i> (L.) G.L. Nesom	Smooth blue aster	D	FU-U	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum laeve</i> (L.) Á. Löve & D. Löve	White panicle aster	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum lanceolatum</i> (Willd.) G.L. Nesom	Calico aster	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum lateriflorum</i> (L.) Á. Löve & D. Löve	Lowrie's blue wood aster	D	.	MD-NJ-PA-VA-WV
	<i>Symphytotrichum lowrianum</i> (Porter) G.L. Nesom	New England aster	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum novae-angliae</i> (L.) G.L. Nesom	New York aster	D	O-FW	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum novi-belgii</i> (L.) G.L. Nesom	Aromatic aster	D	.	MD-NJ-PA-VA-WV
	<i>Symphytotrichum oblongifolium</i> (Nutt.) G.L. Nesom	Bottomland aster	D	F	DE-WV
	<i>Symphytotrichum ontariensis</i> (Wiegand) G.L. Nesom	Late purple aster	D	.	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum patens</i> (Aiton) G.L. Nesom	Thinleaf late purple aster	D	.	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum phlogifolium</i> (Muhl. ex Willd.) G.L. Nesom	Hairy white oldfield aster	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum pilosum</i> (Willd.) G.L. Nesom	Willowleaf aster	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum praealtum</i> (Poir.) G.L. Nesom	Barrens silky aster	D	.	VA
	<i>Symphytotrichum pratense</i> (Raf.) G.L. Nesom	Crookedstem aster	D	F	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum prenanthoides</i> (Muhl. ex Willd.) G.L. Nesom				

Family	Scientific Name	Common Name	Group	Habitat	Range
Brassicaceae	<i>Symphytotrichum puniceum</i> (L.) Á. Löve & D. Löve	Purplestem aster	D	O	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum racemosum</i> (Elliott) G.L. Nesom	Smooth white oldfield aster	D	FW	MD-NJ-PA-VA-WV
	<i>Symphytotrichum shortii</i> (Lindl.) G.L. Nesom	Short's aster	D	.	MD-PA-VA-WV
	<i>Symphytotrichum subulatum</i> (Michx.) G.L. Nesom	Eastern annual saltmarsh aster	D	O	DE-MD-NJ-PA-VA
	<i>Symphytotrichum tenuifolium</i> (L.) G.L. Nesom	Perennial saltmarsh aster	D	O	DE-MD-NJ-VA
	<i>Symphytotrichum undulatum</i> (L.) G.L. Nesom	Wavyleaf aster	D	.	DE-MD-NJ-PA-VA-WV
	<i>Symphytotrichum urophyllum</i> (Lindl.) G.L. Nesom	White arrowleaf aster	D	.	MD-NJ-PA-VA-WV
	<i>Vernonia gigantea</i> (Walter) Trel.	Giant ironweed	D	F	DE-MD-PA-VA-WV
	<i>Vernonia glauca</i> (L.) Willd.	Broadleaf ironweed	D	.	DE-MD-NJ-PA-VA-WV
	<i>Vernonia noveboracensis</i> (L.) Michx.	New York ironweed	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Arabis alpina</i> L.	Alpine rockcress	D	F	VA
	<i>Arabis canadensis</i> L.	Sicklepod	D	.	DE-MD-NJ-PA-VA-WV
	<i>Arabis drummondii</i> A. Gray	Drummond's rockcress	D	FU	DE-NJ
	<i>Arabis glabra</i> (L.) Bernh.	Tower rockcress	D	.	DE-MD-NJ-PA-VA-WV
	<i>Arabis hirsuta</i> (L.) Scop.	Hairy rockcress	D	.	MD-NJ-PA-VA-WV
	<i>Arabis laevigata</i> (Muhl. ex Willd.) Poir.	Smooth rockcress	D	.	DE-MD-NJ-PA-VA-WV
	<i>Arabis lyrata</i> L.	Lyrate rockcress	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Arabis missouriensis</i> Greene	Green rockcress	D	.	PA-MD
	<i>Arabis patens</i> Sull.	Spreading rockcress	D	.	MD-PA-VA-WV
	<i>Arabis serotina</i> Steele	Shale barren rockcress	D	.	VA-WV
	<i>Arabis shortii</i> (Fernald) Gleason	Short's rockcress	D	FU	MD-PA-VA-WV
Cardamine	<i>Cardamine angustata</i> O.E. Schulz	Slender toothwort	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Cardamine bulbosa</i> (Schreb. ex Muhl.) Britton, Sterns & Poggenb.	Bulbous bittercress	D	O	DE-MD-NJ-PA-VA-WV
	<i>Cardamine clematitis</i> Shuttlw. ex A. Gray	Small mountain bittercress	D	O-F	VA
	<i>Cardamine concatenata</i> (Michx.) Sw.	Cutleaf toothwort	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Cardamine diphylla</i> (Michx.) Alph. Wood	Crinkleroot	D	FU-U	DE-MD-NJ-PA-VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
Campanulaceae	<i>Cardamine dissecta</i> (Leavenworth) Al-Shehbaz	Forkleaf toothwort	D	.	VA-WV
	<i>Cardamine douglassii</i> Britton	Limestone bittercress	D	O-FW	MD-NJ-PA-VA-WV
	<i>Cardamine flagellifera</i> O.E. Schulz	Blue Ridge bittercress	D	FU	VA-WV
	<i>Cardamine longii</i> Fernald	Long's bittercress	D	O	DE-MD-NJ-VA
	<i>Cardamine maxima</i> (Nutt.) Alph. Wood	Large toothwort	D	.	NJ-PA-WV
	<i>Cardamine micranthera</i> Rollins	Streambank bittercress	D	O	VA
	<i>Cardamine parviflora</i> L.	Sand bittercress	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Cardamine pennsylvanica</i> Muhl. ex Willd.	Pennsylvania bittercress	D	O-FW	DE-MD-NJ-PA-VA-WV
	<i>Cardamine pratensis</i> L.	Cuckoo flower	D	.	MD-NJ-PA-VA-WV
	<i>Cardamine rotundifolia</i> Michx.	American bittercress	D	O	DE-MD-NJ-PA-VA-WV
	<i>Campanula aparinoides</i> Pursh	Marsh bellflower	D	O	DE-MD-NJ-PA-VA-WV
	<i>Campanula divaricata</i> Michx.	Small bonny bellflower	D	.	MD-VA-WV
	<i>Campanula rotundifolia</i> L.	Bluebell bellflower	D	FU	MD-NJ-PA-VA-WV
	<i>Triodanis biflora</i> (Ruiz & Pav.) Greene	Small Venus' looking-glass	D	.	PA-VA
	<i>Triodanis perfoliata</i> (L.) Nieuwl.	Clasping Venus' looking-glass	D	F-FU	DE-MD-NJ-PA-VA-WV
Convolvulaceae	<i>Ipomoea hederifolia</i> L.	Scarlet creeper	D	FW-F	VA
Comaceae	<i>Ipomoea imperati</i> (Vahl) Griseb.	Beach morning-glory	D	FU	PA
	<i>Ipomoea lacunosa</i> L.	Whitestar	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Ipomoea nil</i> (L.) Roth	Whiteedge morning-glory	D	.	MD-VA
	<i>Ipomoea pandurata</i> (L.) G. Mey.	Man of the earth	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Ipomoea pes-caprae</i> (L.) R. Br.	Bayhops	D	F	PA
	<i>Cornus alternifolia</i> L. f.	Alternatleaf dogwood	ST	F	DE-MD-NJ-PA-VA-WV
	<i>Cornus anomum</i> Mill.	Silky dogwood	S	FW	DE-MD-NJ-PA-VA-WV
	<i>Cornus drummondii</i> C.A. Mey.	Roughleaf dogwood	ST	F	PA
	<i>Cornus foemina</i> Mill.	Stiff dogwood	ST	FW	DE-MD-NJ-VA
	<i>Cornus racemosa</i> Lam.	Gray dogwood	S	F	DE-MD-NJ-PA-VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
Ericaceae	<i>Cornus rugosa</i> Lam.	Roundleaf dogwood	ST	.	DE-MD-NJ-PA-VA-WV
	<i>Cornus sericea</i> L.	Redosier dogwood	ST	.	DE-MD-NJ-PA-VA-WV
	<i>Gaylussacia baccata</i> (Wagenh.) K. Koch	Black huckleberry	S	FU	DE-MD-NJ-PA-VA-WV
	<i>Gaylussacia brachycera</i> (Michx.) A. Gray	Box huckleberry	S	.	DE-MD-PA-VA-WV
	<i>Gaylussacia dumosa</i> (Andrews) Torr. & A. Gray	Dwarf huckleberry	S	F	DE-MD-NJ-PA-VA-WV
	<i>Gaylussacia frondosa</i> (L.) Torr. & Gray ex. Torr.	Blue huckleberry	S	F	DE-MD-NJ-PA-VA-WV
	<i>Lyonia ligustrina</i> (L.) DC.	Maleberry	S	FW	DE-MD-NJ-PA-VA-WV
	<i>Lyonia lucida</i> (Lam.) K. Koch	Fetterbush lyonia	S	FW	VA
	<i>Lyonia mariana</i> (L.) D. Don	Piedmont staggerbush	S	F	DE-MD-NJ-PA-VA
	<i>Rhododendron arborescens</i> (Pursh) Torr.	Smooth azalea	S	FW-F	MD-PA-VA-WV
	<i>Rhododendron atlanticum</i> (Ashe) Rehder	Dwarf azalea	S	F	DE-MD-NJ-PA-VA
	<i>Rhododendron calendulaceum</i> (Michx.) Torr.	Flame azalea	S	.	MD-PA-VA-WV
	<i>Rhododendron canadense</i> (L.) Torr.	Rhodora	S	FW	NJ-PA
	<i>Rhododendron canescens</i> (Michx.) Sweet	Mountain azalea	S	FW	DE-MD-PA
	<i>Rhododendron catawbiense</i> Michx.	Catawba rosebay	S	FU	VA-WV
	<i>Rhododendron cumberlandense</i> E.L. Braun	Cumberland rhododendron	S	.	VA
	<i>Rhododendron maximum</i> L.	Great laurel	ST	F	DE-MD-NJ-PA-VA-WV
	<i>Rhododendron periclymenoides</i> (Michx.) Shinnery	Pink azalea	S	F	DE-MD-NJ-PA-VA-WV
	<i>Rhododendron prinophyllum</i> (Small) Millais	Early azalea	S	F	MD-NJ-PA-VA-WV
	<i>Rhododendron viscosum</i> (L.) Torr.	Swamp azalea	S	O-FW	DE-MD-NJ-PA-VA-WV
	<i>Vaccinium angustifolium</i> Aiton	Lowbush blueberry	S	FU	DE-MD-NJ-PA-VA-WV
	<i>Vaccinium arboreum</i> Marshall	Farkleberry	ST	FU	VA
	<i>Vaccinium caesariense</i> Mack.	New Jersey blueberry	S	O	MD-NJ-PA-VA
	<i>Vaccinium corymbosum</i> L.	Highbush blueberry	S	FW	DE-MD-NJ-PA-VA-WV
	<i>Vaccinium crassifolium</i> Andrews	Creeping blueberry	S	F	VA
	<i>Vaccinium elliptiotii</i> Chapm.	Elliott's blueberry	S	FW	VA

Family	Scientific Name	Common Name	Group	Habitat	Range
Fabaceae	<i>Vaccinium erythrocarpum</i> Michx.	Southern mountain cranberry	S	F	VA-WV
	<i>Vaccinium formosum</i> Andrews	Southern blueberry	S	O-F	MD-NJ-VA
	<i>Vaccinium fuscatum</i> Aiton	Black highbush blueberry	S	FW-F	DE-MD-NJ-PA-VA
	<i>Vaccinium macrocarpon</i> Aiton	Cranberry	S	O	DE-MD-NJ-PA-VA-WV
	<i>Vaccinium myrtilloides</i> Michx.	Velvetleaf huckleberry	S	F-FW	MD-PA-VA-WV
	<i>Vaccinium oxycoccos</i> L.	Small cranberry	S	O	MD-NJ-PA-VA-WV
	<i>Vaccinium pallidum</i> Aiton	Blue Ridge blueberry	S	.	DE-MD-NJ-PA-VA-WV
	<i>Vaccinium stamineum</i> L.	Deerberry	S	FU	DE-MD-NJ-PA-VA-WV
	<i>Vaccinium tenellum</i> Aiton	Small black blueberry	S	FU-U	VA
	<i>Cercis canadensis</i> L.	Eastern redbud	ST	FU-U	DE-MD-NJ-PA-VA-WV
Geraniaceae	<i>Galactia regularis</i> (L.) Britton, Sterns & Poggenb.	Eastern milkpea	D	.	DE-MD-NJ-PA-VA
	<i>Galactia volubilis</i> (L.) Britton	Downy milkpea	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Strophostyles helvola</i> (L.) Elliott	Amberique-bean	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Strophostyles leiosperma</i> (Torr. & A. Gray) Piper	Slickseed fuzzybean	D	.	MD-NJ-PA-VA
	<i>Strophostyles umbellata</i> (Muhl. ex Willd.) Britton	Pink fuzzybean	D	F-FU	DE-MD-NJ-PA-VA
Geraniaceae	<i>Geranium bicknellii</i> Britton	Bicknell's cranesbill	D	.	NJ-PA-VA-WV
	<i>Geranium carolinianum</i> L.	Carolina geranium	D	.	DE-MD-NJ-PA-VA-WV
Hydrophyllaceae	<i>Geranium maculatum</i> L.	Spotted geranium	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Hydrophyllum appendiculatum</i> Michx.	Great waterleaf	D	.	PA-WV
	<i>Hydrophyllum canadense</i> L.	Bluntleaf waterleaf	D	FW-FU	MD-NJ-PA-VA-WV
	<i>Hydrophyllum macrophyllum</i> Nutt.	Largeleaf waterleaf	D	.	MD-PA-VA-WV
	<i>Hydrophyllum virginianum</i> L.	Eastern waterleaf	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Nemophila aphylla</i> (L.) Brummitt	Smallflower baby blue eyes	D	FW	DE-MD-VA-WV
	<i>Phacelia bipinnatifida</i> Michx.	Fernleaf phacelia	D	.	MD-NJ-PA-VA-WV
	<i>Phacelia covillei</i> S. Watson	Coville's phacelia	D	FW	MD-VA
	<i>Phacelia dubia</i> (L.) Trel.	Smallflower phacelia	D	.	DE-MD-PA-VA-WV
			D	.	

Family	Scientific Name	Common Name	Group	Habitat	Range
Lamiaceae	<i>Phacelia fimbriata</i> Michx.	Fringed phacelia	D	.	VA
	<i>Phacelia hirsuta</i> Nutt.	Fuzzy phacelia	D	.	PA
	<i>Phacelia purshii</i> Buckley	Miami mist	D	.	MD-NJ-PA-VA-WV
	<i>Phacelia ranunculacea</i> (Nutt.) Constance	Oceanblue phacelia	D	FW	WV
	<i>Monarda bradburiana</i> Beck	Eastern beebalm	D	.	VA
	<i>Monarda clinopodia</i> L.	White bergamot	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Monarda didyma</i> L.	Scarlet beebalm	D	F	DE-MD-NJ-PA-VA-WV
	<i>Monarda fistulosa</i> L.	Wild bergamot	D	FU-U	DE-MD-NJ-PA-VA-WV
	<i>Monarda media</i> Willd.	Purple bergamot	D	.	DE-MD-NJ-PA-VA-WV
	<i>Monarda punctata</i> L.	Spotted beebalm	D	FU-U	DE-MD-NJ-PA-VA
Liliaceae	<i>Erythronium albidum</i> Nutt.	White fawnlily	M	FU-U	MD-NJ-PA-VA-WV
	<i>Erythronium americanum</i> Ker Gawl.	Dogtooth violet	M	.	DE-MD-NJ-PA-VA-WV
	<i>Erythronium umbilicatum</i> Parks & Hardin	Dimpled troutlily	M	F	MD-VA-WV
	<i>Uvularia grandiflora</i> Sm.	Largeflower bellwort	M	.	MD-PA-VA-WV
	<i>Uvularia perfoliata</i> L.	Perfoliate bellwort	M	FU	DE-MD-NJ-PA-VA-WV
	<i>Uvularia puberula</i> Michx.	Mountain bellwort	M	F-FU	MD-NJ-PA-VA-WV
Malvaceae	<i>Uvularia sessilifolia</i> L.	Sessileleaf bellwort	M	F	DE-MD-NJ-PA-VA-WV
	<i>Hibiscus coccineus</i> Walter	Scarlet rosemallow	D	O	VA
	<i>Hibiscus laevis</i> All.	Halberdleaf rosemallow	D	O	MD-PA-VA-WV
	<i>Hibiscus moscheutos</i> L.	crimsoneyed rosemallow	D	O	DE-MD-NJ-PA-VA-WV
Onagraceae	<i>Oenothera argillicola</i> Mack.	Shale barren evening primrose	D	.	MD-PA-VA-WV
	<i>Oenothera biennis</i> L.	Common evening primrose	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Oenothera clelandii</i> W. Diettr., P.H. Raven & W.L. Wagner	Cleland's evening primrose	D	.	NJ-WV
	<i>Oenothera fruticosa</i> L.	Narrowleaf evening primrose	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Oenothera grandiflora</i> L'Hér. ex Aiton	Largeflower evening primrose	D	.	NJ-PA-WV
	<i>Oenothera grandis</i> (Britton) Smyth	Showy evening primrose	D	.	MD-NJ

Family	Scientific Name	Common Name	Group	Habitat	Range
	<i>Oenothera humifusa</i> Nutt.	Seabeach evening primrose	D	.	DE-MD-NJ-PA-VA
	<i>Oenothera laciniata</i> Hill	Cutleaf evening primrose	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Oenothera linifolia</i> Nutt.	Threadleaf evening primrose	D	.	VA
	<i>Oenothera nutans</i> Atk. & Bartlett	Nodding evening primrose	D	.	MD-NJ-PA-VA-WV
	<i>Oenothera oakesiana</i> (A. Gray) J.W. Robbins ex S. Watson & J.M. Coult.	Oakes' evening primrose	D	.	DE-MD-NJ-PA-VA
	<i>Oenothera parviflora</i> L.	Northern evening primrose	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Oenothera perennis</i> L.	Little evening primrose	D	F	DE-MD-NJ-PA-VA-WV
	<i>Oenothera pilosella</i> Raf.	Meadow evening primrose	D	F-FU	NJ-PA-VA-WV
	<i>Oenothera speciosa</i> Nutt.	Pinkladies	D	.	PA-VA-WV
	<i>Oenothera triloba</i> Nutt.	Stemless evening primrose	D	.	PA-VA
	<i>Oenothera villosa</i> Thunb.	Hairy evening primrose	D	F-FU	MD-NJ-PA-VA-WV
Pontederiaceae	<i>Pontederia cordata</i> L.	Pickersweed	M	O	DE-MD-NJ-PA-VA-WV
Polemoniaceae	<i>Polemonium micranthum</i> Benth.	Annual polemonium	D	FU	PA
Portulacaceae	<i>Polemonium reptans</i> L.	Greek valerian	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Claytonia caroliniana</i> Michx.	Carolina springbeauty	D	FU	MD-PA-VA-WV
	<i>Claytonia rubra</i> (Howell) Tidestr.	Redstem springbeauty	D	.	VA
Primulaceae	<i>Claytonia virginica</i> L.	Virginia springbeauty	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Lysimachia ciliata</i> L.	Fringed loosestrife	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Lysimachia hybrida</i> Michx.	Lowland yellow loosestrife	D	O	DE-MD-NJ-PA-VA-WV
	<i>Lysimachia lanceolata</i> Walter	Lanceleaf loosestrife	D	F	MD-NJ-PA-VA-WV
	<i>Lysimachia quadriflora</i> Sims	Fourflower yellow loosestrife	D	O-FW	MD-PA-VA-WV
	<i>Lysimachia quadrifolia</i> L.	Whorled yellow loosestrife	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Lysimachia radicans</i> Hook.	Trailing yellow loosestrife	D	O	VA
	<i>Lysimachia terrestris</i> (L.) Britton, Sterns & Poggenb.	Earth loosestrife	D	O	DE-MD-NJ-PA-VA-WV
	<i>Lysimachia thysiflora</i> L.	Tufted loosestrife	D	O	MD-NJ-PA-WV
	<i>Lysimachia tonsa</i> (Alph. Wood) Alph. Wood ex Pax & R. Knuth	Southern yellow loosestrife	D	.	VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
Rhamnaceae	<i>Ceanothus americanus</i> L.	New Jersey tea	S	.	DE-MD-NJ-PA-VA-WV
	<i>Ceanothus herbaceus</i> Raf.	Jersey tea	S	.	VA-WV
	<i>Fragaria ×ananassa</i> (Weston) Duchesne ex Rozier (pro sp.)	Strawberry	D	.	NJ-PA-VA-WV
	<i>Fragaria vesca</i> L.	Woodland strawberry	D	FU-U	DE-MD-NJ-PA-VA-WV
	<i>Potentilla virginiana</i> Duchesne	Virginia strawberry	D	FU	DE-MD-NJ-PA-VA-WV
Rosaceae	<i>Potentilla arguta</i> Pursh	Tall cinquefoil	D	.	MD-NJ-PA-VA-WV
	<i>Potentilla canadensis</i> L.	Dwarf cinquefoil	D	.	DE-MD-NJ-PA-VA-WV
	<i>Potentilla norvegica</i> L.	Norwegian cinquefoil	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Potentilla paradoxa</i> Nutt.	Paradox cinquefoil	D	.	PA
	<i>Potentilla rivalis</i> Nutt.	Brook cinquefoil	D	FW	MD-VA
	<i>Potentilla simplex</i> Michx.	Common cinquefoil	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Waldsteinia fragarioides</i> (Michx.) Tratt	Appalachian barren strawberry	D	.	MD-NJ-PA-VA-WV
	<i>Houstonia caerulea</i> L.	Azure bluet	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Houstonia canadensis</i> Willd. ex Roem. & Schult.	Canadian summer bluet	D	.	PA-VA-WV
	<i>Houstonia longifolia</i> Gaertn.	Longleaf summer bluet	D	.	MD-NJ-PA-VA-WV
Salicaceae	<i>Houstonia purpurea</i> L.	Venus' pride	D	.	DE-MD-NJ-PA-VA-WV
	<i>Houstonia pusilla</i> Schoepf	Tiny bluet	D	F-FU	DE-MD-VA
	<i>Houstonia serpyllifolia</i> Michx.	Thymeleaf bluet	D	FW-F	MD-PA-VA-WV
	<i>Salix amygdaloides</i> Andersson	Peachleaf willow	ST	FW	PA
	<i>Salix bebbiana</i> Sarg.	Bebb willow	ST	FW	MD-NJ-PA
	<i>Salix candida</i> Flueggé ex Willd.	Sageleaf willow	S	O	NJ-PA
	<i>Salix caroliniana</i> Michx.	Coastal plain willow	T	O	DE-MD-NJ-PA-VA-WV
	<i>Salix cordata</i> Michx.	Heartleaf willow	S	F	MD-NJ-PA
	<i>Salix discolor</i> Muhl.	Pussy willow	ST	FW	DE-MD-NJ-PA-VA-WV
	<i>Salix eriocephala</i> Michx.	Missouri River willow	ST	FW	DE-MD-NJ-PA-VA-WV
	<i>Salix humilis</i> Marshall	Prairie willow	S	FU	DE-MD-NJ-PA-VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
Saxifragaceae	<i>Salix interior</i> Rowlee	Sandbar willow	ST	O-FW	DE-MD-NJ-PA-VA-WV
	<i>Salix lucida</i> Muhl.	Shining willow	ST	FW	DE-MD-NJ-PA-VA-WV
	<i>Salix myricoides</i> Muhl.	Bayberry willow	ST	FW	PA
	<i>Salix nigra</i> Marshall	Black willow	T	O	DE-MD-NJ-PA-VA-WV
	<i>Salix pedicellaris</i> Pursh	Bog willow	S	O	NJ-PA
	<i>Salix petiolaris</i> Sm.	Meadow willow	ST	FW	NJ-PA
	<i>Salix sericea</i> Marshall	Silky willow	ST	O	DE-MD-NJ-PA-VA-WV
	<i>Salix serissima</i> (L. H. Bailey) Fernald	Autumn willow	ST	O	NJ-PA
	<i>Heuchera alba</i> Rydb.	White alumroot	D	.	VA-WV
	<i>Heuchera americana</i> L.	American alumroot	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Heuchera longiflora</i> Rydb.	Longflower alumroot	D	.	VA-WV
	<i>Heuchera parviflora</i> Bartlett	Littleflower alumroot	D	.	VA-WV
Scrophulariaceae	<i>Heuchera pubescens</i> Pursh	Downy alumroot	D	.	MD-PA-VA-WV
	<i>Agalinis auriculata</i> (Michx.) S.F. Blake	Earleaf false foxglove	D	.	MD-NJ-PA-VA-WV
	<i>Agalinis fasciculata</i> (Elliott) Raf.	Beach false foxglove	D	F	DE-MD-NJ-PA-VA
	<i>Agalinis linfofolia</i> (Nutt.) Britton	Flaxleaf false foxglove	D	O-FW	DE-MD-VA
	<i>Agalinis maritima</i> (Raf.) Raf.	Saltmarsh false foxglove	D	O-FW	DE-MD-NJ-VA
	<i>Agalinis obtusifolia</i> Raf.	Tenlobe false foxglove	D	F-FU	DE-MD-PA-VA
	<i>Agalinis paupercula</i> (A. Gray) Britton	Smallflower false foxglove	D	FW	NJ-PA-VA
	<i>Agalinis purpurea</i> (L.) Pennell	Purple false foxglove	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Agalinis setacea</i> (J.F. Gmel.) Raf.	Threadleaf false foxglove	D	.	DE-MD-NJ-VA
	<i>Agalinis tenuifolia</i> (Vahl) Raf.	Slenderleaf false foxglove	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Penstemon calycosus</i> Small	Longsepal beardtongue	D	FU-U	MD-NJ-PA-VA
	<i>Penstemon canescens</i> (Britton) Britton	Eastern gray beardtongue	D	.	MD-PA-VA-WV
	<i>Penstemon digitalis</i> Nutt. ex Sims	Foxglove beardtongue	D	F	DE-MD-NJ-PA-VA-WV
	<i>Penstemon hirsutus</i> (L.) Willd.	Hairy beardtongue	D	.	DE-MD-NJ-PA-VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
Solanaceae	<i>Penstemon laevigatus</i> Aiton	Eastern smooth beardtongue	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Penstemon pallidus</i> Small	Pale beardtongue	D	FU	MD-NJ-PA-VA-WV
	<i>Penstemon tubaefflorus</i> Nutt.	White wand beardtongue	D	.	PA
	<i>Physalis angulata</i> L.	Cutleaf groundcherry	D	F-FU	DE-NJ-VA
	<i>Physalis cordata</i> Mill.	Heartleaf groundcherry	D	.	PA
	<i>Physalis grisea</i> (Waterf.) M. Martínez	Strawberry-tomato	D	.	NJ-PA-VA
	<i>Physalis heterophylla</i> Nees	Clammy groundcherry	D	.	DE-MD-NJ-PA-VA-WV
	<i>Physalis longifolia</i> Nutt.	Longleaf groundcherry	D	.	DE-MD-NJ-PA-VA-WV
	<i>Physalis pubescens</i> L.	Husk tomato	D	FU-U	MD-NJ-PA-VA-WV
	<i>Physalis virginiana</i> Mill.	Virginia groundcherry	D	.	DE-MD-NJ-PA-VA-WV
Verbenaceae	<i>Physalis walteri</i> Nutt.	Walter's groundcherry	D	.	VA
	<i>Verbena bracteata</i> Cav. ex Lag. & Rodr.	Bigbract verbena	D	FU-U	MD-NJ-PA-VA-WV
	<i>Verbena hastata</i> L.	Swamp verbena	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Verbena scabra</i> Vahl	Sandpaper vervain	D	FW	MD-VA-WV
	<i>Verbena simplex</i> Lehm.	Narrowleaf vervain	D	.	DE-MD-NJ-PA-VA-WV
	<i>Verbena stricta</i> Vent.	Hoary verbena	D	.	DE-NJ-PA-WV
Violaceae	<i>Verbena urticifolia</i> L.	White vervain	D	F	DE-MD-NJ-PA-VA-WV
	<i>Viola affinis</i> Leconte	Sand violet	D	FW	MD-NJ-PA-VA-WV
	<i>Viola appalachensis</i> Henry	Appalachian violet	D	.	MD-PA-WV
	<i>Viola bicolor</i> Pursh	Field pansy	D	F-FU	DE-MD-NJ-PA-VA-WV
	<i>Viola blanda</i> Willd.	Sweet white violet	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Viola brittoniana</i> Pollard	Northern coastal violet	D	F	DE-MD-NJ-PA-VA
	<i>Viola canadensis</i> L.	Canadian white violet	D	F	MD-NJ-PA-VA-WV
	<i>Viola cucullata</i> Aiton	Marsh blue violet	D	O-FW	DE-MD-NJ-PA-VA-WV
	<i>Viola hastata</i> Michx.	Halberdleaf yellow violet	D	FU-U	MD-PA-VA-WV
	<i>Viola hirsutula</i> Brainerd	Southern woodland violet	D	FU	DE-MD-NJ-PA-VA-WV

Family	Scientific Name	Common Name	Group	Habitat	Range
	<i>Viola labradorica</i> Schrank	Alpine violet	D	F	DE-MD-NJ-PA-VA-WV
	<i>Viola lanceolata</i> L.	Bog white violet	D	O	DE-MD-NJ-PA-VA-WV
	<i>Viola mackloskeyi</i> Lloyd	Small white violet	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Viola nephrophylla</i> Greene	Northern bog violet	D	FW	PA-WV
	<i>Viola pedata</i> L.	Birdfoot violet	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Viola pubescens</i> Aiton	Downy yellow violet	D	FU	DE-MD-NJ-PA-VA-WV
	<i>Viola renifolia</i> A. Gray	White violet	D	FW	PA
	<i>Viola rostrata</i> Pursh	Longspur violet	D	FU	MD-NJ-PA-VA-WV
	<i>Viola rotundifolia</i> Michx.	Roundleaf yellow violet	D	F	DE-MD-NJ-PA-VA-WV
	<i>Viola sagittata</i> Aiton	Arrowleaf violet	D	FW-F	DE-MD-NJ-PA-VA-WV
	<i>Viola selkirkii</i> Pursh ex Goldie	Selkirk's violet	D	.	PA
	<i>Viola septentrionalis</i> Greene	Northern woodland violet	D	FU	MD-NJ-PA-VA-WV
	<i>Viola sororia</i> Willd.	Common blue violet	D	F	DE-MD-NJ-PA-VA-WV
	<i>Viola striata</i> Aiton	Striped cream violet	D	FW	DE-MD-NJ-PA-VA-WV
	<i>Viola triloba</i> Schwein.	Three-lobed violet	D	.	MD-NJ-PA-VA-WV
	<i>Viola tripartita</i> Elliott	Three-part violet	D	.	MD-PA-WV
	<i>Viola villosa</i> Walter	Carolina violet	D	FU	VA

THE MARYLAND ENTOMOLOGIST

NOTICE TO CONTRIBUTORS

Contributors should prepare manuscripts according to the following instructions.

Title: The title should be brief, concise, and pertinent.

Abstract: An abstract is required for all long articles; suggested for all biologic studies more than two (2) pages in length; but is not needed for notes, distribution reports, or short observations (especially if two pages or shorter in length). The abstract should provide a capsule description of the main thrust, methods and essential findings of the article. It should contain the scientific name, including species author(s), of the main subject species for most biological studies.

Text: Manuscripts should be submitted in MS Word™ by e-mail or on a CD. Please identify respective file name(s) for text, figure titles, and descriptions of graphs or figures. First mention of a biological organism, in the abstract and text should include the full scientific name, including species author(s). Carefully check the spelling of all scientific names. All genera and species should be in italics. Capitalize “official” common names for faunal species. Short articles and general notes (20 pages or less) are preferred. Longer manuscripts may be assessed page charges. Color “copy ready” illustrations, pictures, or digital images are preferred.

References: References should be given in an author-date format: (Lynch 1987); (Lynch 1987, 1988); (Lynch, in press); (Lynch, in litt.); (Lynch, pers. comm.); (Lynch and Lynch 1989); and (Lynch et al.) for 3 or more authors. Provide evidence of acceptance for works “in press,” or cite as “unpublished”, “in litt.” (written), or “pers. comm.” (verbal); written permission is suggested as well. Citations shall be listed alphabetically, under LITERATURE CITED, as follows: **Articles:** Lynch, J.F. 1987. An annotated checklist and key to the species of ants (Hymenoptera: Formicidae) of the Chesapeake Bay region. *The Maryland Naturalist* 31(3-4):61-105. Do not abbreviate the titles of journals. **Books:** Fisher, B.L., and S.P. Cover. 2007. *Ants of North America: a guide to the genera*. University of California Press, Berkeley and Los Angeles, CA. 194 pp. **Internet:** Entomological Society of America. 2011. Common names of insects database. Available at: <http://entsoc.org/common-names>. Accessed 11 July 2011.

Tables: Tables, graphs and line drawings should be created electronically in black and white. Color should only be used when absolutely necessary for clarity.

Illustrations: Photographs or high-definition images may be accepted if necessary or desired by the author(s) to support the text. Reproduction of photos or images may increase printing costs and authors will be expected to pay any extra charges. Photographs should be submitted in color. Figure numbers, as cited in the text, and figure legends should be keyed to each respective photograph.

The Maryland Entomologist is published annually by the Maryland Entomological Society. There are four numbers per volume. Original articles or reports on geographic or temporal distribution (primarily pertaining to Maryland and the Mid-Atlantic region), ecology, biology, morphology, genetics, systematics, behavior, etc., are welcome. Notes on distribution, behavior, occurrence, migration, life history, and other biological topics will be published. All submissions are subject to editorial review and acceptance. Articles will be peer-reviewed. Send submissions to the Editor: Eugene J. Scarpulla, 14207 Lakerun Court, Bowie, MD 20720-4861, ejscarp@comcast.net.

CONTENTS

Editor’s Note..... 1

Specialist Bees of the Mid-Atlantic: Host Plants and Habitat Conservation
Jarrod Fowler..... 2

COVER PHOTOGRAPH

A female long-horned bee, *Melissodes denticulatus* Smith (Hymenoptera: Apidae), a specialist on New York ironweed, *Vernonia noveboracensis* (L.) Michx. (Asteraceae). Photographed near Lake Elkhorn, Columbia, Howard County, Maryland, 16 August 2015.

Photographed by Richard L. Orr